arXiv:0811.3726v4 [math.GT] 30 Sep 2009

CONFIGURATION SPACE INTEGRALS FOR EMBEDDING SPACES AND THE HAEFLIGER INVARIANT

KEIICHI SAKAI

Abstract. Let $\mathcal{K}_{n,j}$ be the space of long j-knots in $\mathbb{R}^n$. In this paper we introduce a graph complex $\mathcal{D}^*$ and a linear map $I : \mathcal{D}^* \to \Omega^*_{DR}(\mathcal{K}_{n,j})$ via configuration space integral, and prove that (1) when both $n > j \geq 3$ are odd, I is a cochain map if restricted to graphs with at most one loop component, (2) when $n-j \geq 2$ is even, I is a cochain map if restricted to tree graphs, and (3) when $n-j \geq 3$ is odd, I added a correction term produces a $(2n-3j-3)$-cocycle of $\mathcal{K}_{n,j}$ which gives a new formulation of the Haefliger invariant when $n = 6k$, $j = 4k-1$ for some k.

1. Introduction

The aim of this research is to study the topology of the space of long knots.

Definition 1.1. A *long j-knot* in $\mathbb{R}^n$ is an embedding $f : \mathbb{R}^j \hookrightarrow \mathbb{R}^n$ which is standard at infinity:

$$f(x) = (x, 0) \in \mathbb{R}^j \times \{0\}^{n-j}, \quad x \notin [-1, 1]^j.$$

Denote by $\mathcal{K}_{n,j}$ the space of long j-knots in $\mathbb{R}^n$, equipped with C^∞-topology.

In this and the forthcoming papers [16] we will develop the methods to study $\mathcal{K}_{n,j}$ originated in perturbative Chern-Simons theory. This method was used by various authors [1, 3, 10] to give some integral expressions of the finite type invariants for knots in $\mathbb{R}^3$. Cattaneo, Cotta-Ramusino and Longoni [5] generalized their constructions to define a cochain map from a certain graph complex to the de Rham complex of the space of (long) 1-knots in $\mathbb{R}^n$, $n > 3$, via fiber-integrations over configuration spaces. Not only the trivalent graphs correspond to finite type invariants, but non-trivalent graphs can also work in their framework. Indeed, some non-trivalent graph cocycles produce non-trivial cohomology classes [11, 15].

Another generalization was done by Rossi [14] and Cattaneo-Rossi [6], who proved the invariance of (order two) Bott invariant and order three invariant for long m-knots in $\mathbb{R}^{m+2}$ for $m \geq 2$. Following their work, Watanabe [20] proved that there is one finite type invariant for long "ribbon" m-knots [7] in $\mathbb{R}^{m+2}$, $m \geq 3$ odd, at each even order. These invariants also come from trivalent graphs via the perturbative method.

In this paper we introduce graph complexes in the same manner as [5] generated by more general graphs than those in [6, 14, 20]. We prove that some graph cocycles produce cohomology classes of $\mathcal{K}_{n,j}$ via configuration space integrals.

Date: September 30, 2009.

2000 *Mathematics Subject Classification.* Primary 58D10; Secondary 57Q45, 81Q30.

Key words and phrases. The space of long j-knots; Graph complex; Configuration space integral; Haefliger invariant.

Theorem 1.2. *There exist graph complexes*[1] $\mathcal{D}_g^{k,*}$, $k \geq 1$ *and* $g \geq 0$, *spanned by graphs of first Betti number* g *(after its 'small loops' are removed) and linear maps* $I : \mathcal{D}_g^{k,l} \to \Omega_{DR}^{k(n-j-2)+(g-1)(j-1)+l}(\mathcal{K}_{n,j})$ *given by configuration space integral. The map* I *is a cochain map when* $n-j \geq 2$ *is even and* $g = 0$, *or both* $n > j \geq 3$ *are odd and* $g = 1$.

Unfortunately it is not known how 'big' the graph cohomology $H^*(\mathcal{D}_0^*)$ is. But in [16] we will prove that $H^*(\mathcal{D}_1^*)$ is not trivial, and the map I produces non-trivial cohomology classes of $\mathcal{K}_{n,j}$ and $\overline{\mathcal{K}}_{n,j}$ for various n and j, which generalize the 'finite type invariants' for 'long ribbon knots' [7, 20]. Here $\overline{\mathcal{K}}_{n,j}$ is the space of long j-knots 'modulo immersions' (see for example [16, 17]). When $n-j$ is odd, we have no idea to prove that I is a cochain map, since we cannot ignore the contributions of some 'hidden faces' of the boundary of configuration spaces. So we will need some correction terms (different from that given in Theorem 1.3). See [16].

The graph complex introduced in [5] is conjectured to give all the real cohomology classes of $\mathcal{K}_{n,1}$, while it is not clear whether the graphs appeared in [6, 14, 20] are enough for $H_{DR}^*(\mathcal{K}_{n,j})$. But our graph complex $\mathcal{D}^* = \bigoplus_g \mathcal{D}_g^*$ contains more general graphs, so we might be able to conjecture that our graph complex would describe whole $H_{DR}^*(\mathcal{K}_{n,j})$. The next Theorem 1.3 confirms the conjecture; via perturbative method we can give a new formulation of the *Haefliger invariant* [8, 9].

Theorem 1.3. *Suppose* $n-j \geq 3$ *is odd and* $2n-3j-3 \geq 0$. *Then a graph cocycle* $H \in \mathcal{D}_0^{2,0}$ *produces a non-trivial cohomology class* $\mathcal{H} := [I(H) + c] \in H_{DR}^{2n-3j-3}(\mathcal{K}_{n,j})$, *where* c *is some correction term. Moreover when* $n = 6k$ *and* $j = 4k-1$, $\mathcal{H} \in H_{DR}^0(\mathcal{K}_{6k,4k-1})$ *is nothing but the Haefliger invariant (up to sign).*

The correction term c is added to kill some contribution of the 'anomalous face' of a compactified configuration space. This contribution is an obstruction for I to be a cochain map. See §4 and §5 for details.

The Haefliger invariant, an isotopy invariant for (long) $(4k-1)$-knots in $6k$-space, was originally defined by using a $4k$-manifold bounded by the knot in $(6k+1)$-space. Instead of such additional data, we use the generators of cohomology of configuration space.

This paper is organized as follows. We define the graph complexes in §2 and describe the integration map I in detail in §3. Several vanishing results are proved in §5 to prove Theorem 1.2. The class $\mathcal{H} \in H_{DR}^{2n-3j-3}(\mathcal{K}_{n,j})$ is studied in §4. To prove $d\mathcal{H} = 0$ we need some of the results in §5, which hold even if $n-j$ is odd.

Acknowledgment. The author expresses his great appreciation to Professor Toshitake Kohno for his encouragement, to Tadayuki Watanabe for many useful suggestions and comments, and to Masamichi Takase for his idea to improve Theorem 1.3. The author is partially supported by the Grant-in-Aid for Young Scientists (B), MEXT, Japan, by The Sumitomo Foundation, and by The Iwanami Fujukai Foundation.

2. GRAPH COMPLEXES

This and the next sections provide preliminaries for not only the subsequent sections but the forthcoming paper [16]. In this section we introduce more general graphs than those in [6, 14, 20], including 'degenerate graphs.'

---

[1] The combinatorial meaning of the number k will be specified in Definition 2.2.

2.1. **Graphs.** The *graphs* appearing here have two types of vertices. One is the *external* vertex (or shortly *e-vertex*), which is depicted by $\circ$, while the other is the *internal* vertex (shortly *i-vertex*), depicted by $\bullet$.

Similarly there are two types of edges. One is θ*-edge*, which is depicted by a dotted line, and another is η*-edge*, depicted by a solid line. We suppose that all the endpoints of η-edges are i-vertices.

A θ-edge e can form a *small loop* at an i-vertex p, that is, e may have exactly one i-vertex p as its endpoint. A single η-edge is not allowed to be a small loop, but an η-edge together with a θ-edge can form a loop at an i-vertex, called a *double loop*. When we count the edges of a graph, we count a double loop twice, regarding it as consisting of an η-loop and a θ-loop. But a double loop raises the first Betti number of the graph by one.

Definition 2.1. A vertex v of a graph is *admissible* if it is

(1) an i-vertex of valence ≥ 1 with at least one θ-edge (possibly a loop) emanating from v, or
(2) an e-vertex of valence ≥ 3, with only θ-edges (which are not loops) emanating from v

(see Figure 2.1). A graph is called *admissible* if all its vertices are admissible.



FIGURE 2.1. Admissible vertices

Figure 2.2 shows an example of an admissible graph. There might be an i-vertex which is adjacent to more than one θ-edges, an e-vertex of valency ≥ 4, and so on. Such vertices did not appear in [6, 14, 20].



FIGURE 2.2. An example of an admissible graph for odd n and j

In §3 the graphs will be regarded as in $\mathbb{R}^n$ (Figure 2.3); the set of vertices will become a configuration of points in $\mathbb{R}^n$, where the i-vertices are on $\mathbb{R}^j$ (or a long j-knot) embedded in $\mathbb{R}^n$. Each edge will correspond to a 'direction map' determined by its endpoints, or to the volume form of sphere pulled back by the direction map. The η-edges correspond to directions in $\mathbb{R}^j$, while θ-edges to those in $\mathbb{R}^n$.

Afterward we will need to fix the orientations of configuration spaces and the signs of the volume forms. For these purposes we 'decorate' the graphs as follows (see Figure 2.4).



FIGURE 2.3. The graph from Figure 2.2 in $\mathbb{R}^n$

(1) The i-vertices are labeled by $1, \dots, s$ and the e-vertices are labeled by $s+1, \dots, s+t$ for some suitable s and t.
(2) When both n and j are odd, all the edges are oriented, all the loops are ordered and each loop is given the sign ± 1.
(3) When n is odd and j is even, all the θ-edges are oriented, while all the η-edges are labeled.
(4) When n is even and j is odd, all the θ-edges (including those in double loops) are labeled, while all the η-edges are oriented. The double loops are given other labels $1, 2, \dots$ than those for all the θ-edges. A double loop is given a sign ± 1.
(5) When both n and j are even, all the edges (including those in double loops) are labeled. The small / double loops are given other labels.



FIGURE 2.4. Decorations of graphs

2.2. **Space of graphs.** Below we assume that all the graphs are admissible and decorated unless otherwise stated.

Definition 2.2. For a graph Γ, define

$$\operatorname{ord}\Gamma := \sharp\{\theta\text{-edges of } \Gamma\} - \sharp\{\text{e-vertices of } \Gamma\},$$
$$\deg\Gamma := 2\sharp\{\theta\text{-edges of } \Gamma\} - 3\sharp\{\text{e-vertices of } \Gamma\} - \sharp\{\text{i-vertices of } \Gamma\}$$

(for example, $\operatorname{ord}\Gamma = 7$ and $\deg\Gamma = 5$ for the graph Γ in Figure 2.2; see below for more explanations). We denote by $\mathcal{D}^{k,l}$ the vector space spanned by admissible decorated graphs with ord $= k$, deg $= l$ modulo the subspace spanned by

$$\Gamma' - (-1)^{j\operatorname{sign}\sigma + n\operatorname{sign}\tau + a + b + \operatorname{sign}\rho}\Gamma \quad \text{and} \quad \Gamma'',$$

where Γ' is obtained from Γ by

- permuting the labels of i- and e-vertices of Γ by $\sigma \in \mathfrak{S}_s$ and $\tau \in \mathfrak{S}_t$ respectively (s and t are the numbers of i- and e-vertices of Γ respectively),
- reversing a oriented edges of Γ, and
- switching b signs and permuting the labels of small / double loops by ρ,

and Γ'' is a graph with 'multiple η- (or θ-) edges,' i.e., there are two vertices p, q which are joined by two or more η- (or θ-) edges (we allow p, q joined by two edges, one η-edge and one θ-edge). Moreover we introduce one more relation; $\Gamma \sim 0$ in $\mathcal{D}^*$ if $n-j$ is odd and Γ is a graph with at least one small loop, or if n is odd and Γ has a double loop.

Remark 2.3. The sign $j\operatorname{sign}\sigma + n\operatorname{sign}\tau$ will correspond to the orientation sign of the configuration space. In §3 we will associate the volume forms of spheres of even dimensions with the oriented edges. Reversing an oriented edge corresponds to pull-back via the antipodal map, hence yields a sign -1. Unoriented edges correspond to differential forms of odd degrees, so permuting the labels yields a sign.

The meaning of $\operatorname{ord}\Gamma$ is as follows. In Figure 2.3, if we contract $\mathbb{R}^j$ together with η-edges regarded as in $\mathbb{R}^j$, then we obtain a one dimensional CW complex whose edges are θ-edges. Its first Betti number is equal to $\operatorname{ord}\Gamma$.

To explain the meaning of $\deg\Gamma$, we need some terminologies.

Definition 2.4. An admissible vertex is said to be *non-degenerate* if it is

- an i-vertex with exactly one θ-edge (and possibly many η-edges) emanating from it, or
- a tri-valent e-vertex.

All other vertices are said to be *degenerate.*

For example, all the vertices in Figure 2.1, and the vertices 1, 2, 6, 7 and 9 of the graph in Figure 2.2 are non-degenerate.

Remark 2.5. It can be easily understood that "non-degenerate vertex" is the same notion as "trivalent vertex" in [5]. All the vertices of the graphs appeared in [6, 14, 20] are non-degenerate.

Lemma 2.6. *We have* $\deg\Gamma \geq 0$ *for any admissible graph* Γ*, and* $\deg\Gamma = 0$ *if and only if all the vertices of* Γ *are non-degenerate.*

Proof. This Lemma is obvious by the definition of admissible vertices; at least one θ-edge emanates from any i-vertex of a graph and at least three θ-edges emanate from any i-vertex. This implies

$$2\sharp\{\theta\text{-edges}\} \geq 3\sharp\{\text{e-vertices}\} + \sharp\{\text{i-vertices}\}$$

and the equality holds if and only if exactly one (resp. three) θ-edge emanates from any i-vertices (resp. e-vertices), that is, all the vertices are non-degenerate. □

Remark 2.7. Let Γ be an admissible graph of $\operatorname{ord}\Gamma = k$, $\deg\Gamma = l$. Then Γ has $2k - l$ vertices; it is a direct consequence of the definition. In particular, if Γ is non-degenerate ($l = 0$), then the number of all vertices is $2k$. In [20] the half of the number of the vertices of a (non-degenerate) graph is called its 'degree.' Thus our terminology 'ord' is a generalization of the 'degree' in [20].

2.3. **Coboundary operation.**

Definition 2.8. Let Γ be a graph and $e = \overrightarrow{pq}$ its (possibly oriented) edge (but not a loop). Define a new graph Γ/e as follows (see Figure 2.5).

(1) When e is an η-edge (then endpoints p, q are both internal), define Γ/e by contracting e, that is, identifying the endpoints p, q of e and removing the edge e. The decoration of Γ/e is derived from that of Γ; the vertex of Γ/e where the contraction occurred is re-labeled by $\min\{p, q\}$, and all the labels of the vertices of Γ bigger than $\max\{p, q\}$ are decreased by one. The labels of the other vertices remain unchanged. When j is even and e is the i-th edge, then the labels of other edges bigger than i is decreased by one.

(2) When e is a θ-edge and at least one of p, q is an e-vertex, then Γ/e is defined in the same way as above. If both p, q are external, then the vertex where the contraction occurred is also external. If one of p, q is internal, then the resulting vertex is internal.

(3) When e is a θ-edge with both p, q being i-vertices, then Γ/e is obtained from Γ by identifying the vertices p and q, but not removing the edge e. The edge e becomes a small loop at the i-vertex $\min\{p, q\}$. The labeling of Γ/e is determined similarly as above. When n and j are odd, its sign is $+1$ (resp. -1) if $p < q$ (resp. $p > q$). This small loop is labeled by a if Γ has $(a-1)$ small loops.

(4) When e is the η-edge of the multiple edges joining two i-vertices, then Γ/e is obtained from Γ by identifying the vertices p and q, and attaching a double loop at p. This double loop is labeled by a if Γ has $(a-1)$ loops. When j is odd, the sign ± 1 is given similarly as in (3). We do not define Γ/e for a θ-edge of the multiple edges.



FIGURE 2.5. Contractions of edges

We should notice that a graph Γ/e of type (3) in Figure 2.5 is ruled out in $\mathcal{D}^*$ when $n-j$ is odd, and similarly a graph Γ/e of type (4) is ruled out when n is odd (see Definition 2.2).

We would like to define the operator δ by

$$\delta\Gamma = \sum_{e \in E(\Gamma)\setminus\{\text{loops}\}} (-1)^{\tau(e)} \Gamma/e,$$

by giving some suitable signs $\tau(e)$, where $E(\Gamma)$ is the set of edges of Γ.

Proposition 2.9. *If we define the signs $\tau(e)$ as in (1)-(5) below, then the operator δ is well-defined and determines a coboundary operation*

$$\delta : \mathcal{D}^{k,l} \longrightarrow \mathcal{D}^{k,l+1},$$

that is, $\delta \circ \delta = 0$. Thus $\{\mathcal{D}^{k,*}, \delta\}$ is a cochain complex for any k.

(1) *Let both n and j be odd. For any oriented edge $e = \overrightarrow{pq}$, define $\tau(e)$ by*

$$\tau(e) := \begin{cases} q & p < q, \\ p+1 & p > q. \end{cases} \tag{2.1}$$

(2) *Let both n and j be even. For the i-th edge e, define $\tau(e)$ by*

$$\tau(e) := \begin{cases} i & e \text{ is of type (1), (2) in Figure 2.5,} \\ u+1 & e \text{ is of type (3) in Figure 2.5,} \end{cases}$$

where u is the number of small / double loops of Γ.

(3) *Let n be even and j be odd. For any oriented η-edge $e=\overrightarrow{pq}$, define $\tau(e)$ by* (2.1)*. For the i-th θ-edge $e=pq$, $p<q$ with q being an e-vertex, define $\tau(e):=i+s+1$, where s is the number of i-vertices of Γ.*

(4) *Let n be odd and j be even. For the i-th η-edge e, define $\tau(e):=i+t+1$ where t is the number of e-vertices of Γ. For any oriented θ-edge $e=\overrightarrow{pq}$, define $\tau(e)$ by* (2.1)*.*

(5) *Consider the case that n is even, and p and q are i-vertices joined by 'multiple edges,' one η-edge and one θ-edge. If j is odd and the η-edge is oriented from p to q, then τ is given by* (2.1)*. If j is even, then $\tau=u+1$.*

Remark 2.10. The signs in Proposition 2.9 correspond to those of induced orientations of the boundary strata of configuration spaces; see §5.5.

The proof is completely similar to [5, Theorem 4.2]; choose two edges e_1 and e_2, and contract them in two different orders, then we obtain the same graph with opposite signs. Notice that if $n-j$ is odd (resp. n is odd) the case (3) (resp. (4)) in Figure 2.5 does not occur.

3. Configuration space integral

3.1. **Configuration spaces.** For any space M, denote the space of configurations in M by

$$C^o_k(M) := \{(x_1,\dots,x_k)\in M^k \mid x_p\neq x_q \text{ if } p\neq q\}.$$

Let Γ be a (possibly non-admissible) decorated graph with s i-vertices, t e-vertices and u loops. Define the configuration space associated with (vertices of) Γ by

$$C^o_\Gamma := \left\{ \begin{array}{c} (f;x_1,\dots,x_s;y_{s+1},\dots,y_{s+t}) \\ \in \mathcal{K}_{n,j}\times C^o_s(\mathbb{R}^j)\times C^o_t(\mathbb{R}^n) \end{array} \,\middle|\, \begin{array}{c} f(x_p)\neq y_q, \\ \forall p,q \end{array} \right\} \times (S^{j-1})^u.$$

We think of i-vertex p (resp. e-vertex q) as corresponding to $x_p\in\mathbb{R}^j$ (resp. $y_q\in\mathbb{R}^n$) for all $1\le p\le s$ (resp. $s+1\le q\le s+t$). The S^{j-1}-factors will be used to define a differential form ω_e for a loop e (see below). There is a projection

$$\pi_\Gamma : C^o_\Gamma \longrightarrow \mathcal{K}_{n,j}.$$

We will denote its fiber over f by $C^o_\Gamma(f)$.

3.2. **Differential forms associated to graphs.** Let $e=\overrightarrow{pq}$ be an (oriented) edge or a loop of Γ. To e we will assign a differential form $\omega_e\in\Omega^*_{DR}(C^o_\Gamma)$ as follows.

First consider the case that e is not a loop (thus $p\neq q$). When e is an η-edge (then $p,q\le s$), define the 'direction map' $\varphi^\eta_e : C^o_\Gamma\to S^{j-1}$ by

$$\varphi^\eta_e(f,x,y,v) := \frac{x_q-x_p}{|x_q-x_p|}\in S^{j-1}.$$

When e is a θ-edge, define $\varphi^\theta_e : C^o_\Gamma\to S^{n-1}$ by

$$\varphi^\theta_e(f,x,y,v) := \frac{z_q-z_p}{|z_q-z_p|}\in S^{n-1},$$

where

$$z_p=\begin{cases} f(x_p) & \text{if the vertex } p \text{ is internal (thus } p\le s),\\ y_p & \text{if the vertex } p \text{ is external (thus } p>s).\end{cases}$$

Notation. We denote by $vol_{S^{N-1}}$ the *(anti-)symmetric* volume form of S^{N-1} ($N = j$ or n). Namely, $vol_{S^{N-1}}$ is an $(N-1)$-form of S^{N-1} with total integral one and $\iota^* vol_{S^{N-1}} = (-1)^N vol_{S^{N-1}}$ for the antipodal map $\iota : S^{N-1} \to S^{N-1}$.

Define the differential forms $\omega_e \in \Omega^*_{DR}(C^o_\Gamma)$ as $\eta_e := (\varphi^\eta_e)^* vol_{S^{j-1}}$ or $\theta_e := (\varphi^\theta_e)^* vol_{S^{n-1}}$, according to whether e is an η-edge or a θ-edge. Notice that, if e is not oriented, then the map φ_e has ambiguity of signs, but in such a case the corresponding volume form is of odd degree and is invariant under the antipodal map of spheres. Hence the form ω_e is well defined.

When $n-j$ is even, we also assign differential forms to small loops. For the a-th small loop e with sign ε (which is always $+1$ when n is even) at the i-vertex p, define $D_a : C^o_\Gamma \to S^{n-1}$ by

$$D_a(f,x,y,v) := \epsilon \cdot \frac{df_{x_p}(v_a)}{|df_{x_p}(v_a)|},$$

here $df_{x_p} : T_{x_p}\mathbb{R}^j \to T_{f(x_p)}\mathbb{R}^n$ is the derivation map. The differential form associated with e is $\omega_e := D^*_a vol_{S^{n-1}} \in \Omega^{n-1}_{DR}(C^o_\Gamma)$. Such a form is not needed when $n-j$ is odd (see Definition 2.2).

When n is even, to the a-th double loop at the i-vertex p with sign ϵ (which is $+1$ when j is even), we assign a map $\tilde{D}_a : C^o_\Gamma \to S^{j-1} \times S^{n-1}$ defined by

$$\tilde{D}_a(f,x,y,v) := \left(\epsilon v_a, \frac{df_{x_p}(v_a)}{|df_{x_p}(v_a)|} \right)$$

and a differential form $\omega_e := \tilde{D}^*_a(vol_{S^{j-1}} \times vol_{S^{n-1}}) \in \Omega^{n+j-2}_{DR}(C^o_\Gamma)$.

Define the differential form ω_Γ by

$$\omega_\Gamma := \bigwedge_{e \in E(\Gamma)} \omega_e \in \Omega^*_{DR}(C^o_\Gamma),$$

here ω_e's for labeled edges must be ordered according to the labels, since by definition they are odd forms. Since non-labeled edges correspond to even forms, we need not to mention the order of corresponding forms.

3.3. **Fiber integration and compactified configuration spaces.** We would like to define $I(\Gamma) \in \Omega^*_{DR}(\mathcal{K}_{n,j})$ by integrating ω_Γ along the fiber of $\pi_\Gamma : C^o_\Gamma \to \mathcal{K}_{n,j}$. It is not clear whether such an integral converges, because the fiber of π_Γ is not compact. This difficulty has been resolved in [2, 3]; for any manifold M, we can construct a compact manifold $C_k(M)$ with corners so that its interior is $C^o_k(M)$, by 'blowing up' all the diagonals of M^k. We can smoothly extend the direction maps like φ's, the projections $C^o_k(M) \to C^o_{k-l}(M)$ ($l > 0$) and the evaluation map

$$ev : \mathcal{K}_{n,j} \times C^o_k(\mathbb{R}^j) \longrightarrow C^o_k(\mathbb{R}^n), \quad (f; x_1, \dots, x_k) \longmapsto (f(x_1), \dots, f(x_k))$$

onto $C_k(M)$.

Let s and t be the numbers of i- and e-vertices of Γ respectively. Define C'_Γ by the pull-back square

$$\begin{tikzcd}
C'_\Gamma \arrow[r] \arrow[d] & C_{s+t}(\mathbb{R}^n) \arrow[d, "pr_s"] \\
\mathcal{K}_{n,j} \times C_s(\mathbb{R}^j) \arrow[r, "ev"] & C_s(\mathbb{R}^n)
\end{tikzcd}$$

where pr_s is the first s projection, and define $C_\Gamma := C'_\Gamma \times (S^{j-1})^u$ (u is the number of loops of Γ). The differential form ω_Γ is defined on C_Γ since the direction maps φ's are well defined on C_Γ. The natural projection $\pi_\Gamma : C_\Gamma \to \mathcal{K}_{n,j}$ is defined and is a fibration with compact fibers. Thus the integration

$$I(\Gamma) := (\pi_\Gamma)_* \omega_\Gamma \in \Omega^*_{DR}(\mathcal{K}_{n,j})$$

along the fiber is well defined. The degree of $I(\Gamma)$ is given by

$$\begin{aligned}
&\deg \omega_\Gamma - \dim \mathrm{fib}(\pi_\Gamma) \\
&= (n-1)\sharp\{\theta\} + (j-1)\sharp\{\eta\} - j\sharp\{\bullet\} - n\sharp\{\circ\} - (j-1)u \\
&= (n-3)(\sharp\{\theta\} - \sharp\{\circ\}) + (j-1)(\sharp\{\eta\} - \sharp\{\bullet\} - u) + 2\sharp\{\theta\} - 3\sharp\{\circ\} - \sharp\{\bullet\} \\
&= (n-3)\mathrm{ord}\,\Gamma + (j-1)(\sharp\{\eta\} - \sharp\{\bullet\} - u) + \deg \Gamma.
\end{aligned}$$

Suppose that a one-dimensional CW complex $\Gamma \setminus \{\text{small loops}\}$ has g loop components (that is, the first Betti number of it is g). Then

$$\sharp\{\eta\} - \sharp\{\bullet\} - u = -\sharp\{\theta\} + \sharp\{\circ\} + (g-1) = -\mathrm{ord}\,\Gamma + (g-1),$$

and hence

$$\deg I(\Gamma) = (n-j-2)\mathrm{ord}\,\Gamma + (g-1)(j-1) + \deg \Gamma.$$

Proposition 3.1 ([5])**.** *The form $I(\Gamma)$ depends only on the equivalence class of $\Gamma \in \mathcal{D}^*$.*

Proof. This is because the space $\mathcal{D}^*$ is arranged so that the map I is compatible with the permutations of coordinates and the antipodal map of spheres (see [5] for details). But there are two point we should stress here; one is that we have to temporarily gave the labels to loops when j is odd. But the choices of the labeling do not change the form $I(\Gamma)$ since the permutation on the $(S^{j-1})^u$-factor of the fiber $C_\Gamma(f)$ is an orientation preserving diffeomorphism and preserves the form ω_Γ.

Another is that we ruled out the graphs with small loops when $n - j \geq 3$ is odd, and those with double loops when n is odd (see Definition 2.2). This is because the differential forms arising from such graphs are zero (see Lemmas 3.2, 3.3). □

Lemma 3.2. *When $n - j$ is odd, then the differential form $I(\Gamma)$ is zero for any graph Γ with at least one small loop.*

Proof. Let e be a small loop of Γ. Define a fiberwise involution $F : C_\Gamma \to C_\Gamma$ as the antipodal map on S^{j-1} corresponding to e, and the identity on the other factors. The orientation sign of the involution on the fiber of π_Γ is $(-1)^j$, while $F^*\omega_\Gamma = (-1)^n \omega_\Gamma$ since

$$\begin{aligned}
F^*\omega_e &= \left(\frac{df}{|df|} \circ \iota_{S^{j-1}}\right)^* vol_{S^{n-1}} = \left(\iota_{S^{n-1}} \circ \frac{df}{|df|}\right)^* vol_{S^{n-1}} \\
&= (-1)^n \left(\frac{df}{|df|}\right)^* vol_{S^{n-1}} = (-1)^n \omega_e
\end{aligned}$$

and other ω_e's do not change. Thus

$$(\pi_\Gamma)_*\omega_\Gamma = (\pi_\Gamma \circ F)_* F^* \omega_\Gamma = (-1)^{j+n} (\pi_\Gamma)_* \omega_\Gamma = -(\pi_\Gamma)_*\omega_\Gamma$$

since $n + j$ is odd, and hence $(\pi_\Gamma)_*\omega_\Gamma = 0$. □

Lemma 3.3. *When n is odd, then the differential form $I(\Gamma)$ is zero for any graph Γ with at least one double loop.*

Proof. The fiberwise involution F in the proof of Lemma 3.2 has an orientation sign $(-1)^j$ on each fiber, but in the case here $F^*\omega_\Gamma = (-1)^{n+j}\omega_\Gamma$ since F^* affects $vol_{S^{j-1}}$ and $vol_{S^{n-1}}$ as the antipodal map. Thus

$$(\pi_\Gamma)_*\omega_\Gamma = (-1)^{n+2j}(\pi_\Gamma)_*\omega_\Gamma = -(\pi_\Gamma)_*\omega_\Gamma.$$

□

Thus we obtain linear maps

$$I : \mathcal{D}^{k,l}_g \longrightarrow \Omega^{(n-j-2)k+(g-1)(j-1)+l}_{DR}(\mathcal{K}_{n,j}),$$

where $\mathcal{D}^{k,l}_g$ is the subspace of $\mathcal{D}^{k,l}$ spanned by the graphs Γ whose first Betti numbers are g after its small loops (not double loops) are removed. It can be easily seen that $\mathcal{D}^{k,*}_g$ forms a subcomplex of $\mathcal{D}^{k,*}$ for any g (we regard the contraction (4) in Figure 2.5 as preserving the first Betti number).

Here we restate the last half of Theorem 1.2.

Theorem 3.4. *Suppose $n-j \geq 2$ is even and $j \geq 2$. The integration map I : $\mathcal{D}^{k,*}_g \to \Omega^{(n-j-2)k-(j-1)+*}_{DR}(\mathcal{K}_{n,j})$ is a cochain map if (1) both $n > j \geq 2$ are even and $g = 0$, or (2) both $n > j \geq 3$ are odd and $g = 0, 1$.*

Theorem 3.4 is a direct consequence of Theorems 5.11, 5.12, which are proved in similar ways as the results in [5, 14, 20]. The key step is the generalized Stokes theorem (see [5]);

$$dI(\Gamma) = (\pi_\Gamma)_*(d\omega_\Gamma) + (-1)^{\deg\omega_\Gamma+1}(\pi^\partial_\Gamma)_*\omega_\Gamma = (-1)^{\deg\omega_\Gamma+1}(\pi^\partial_\Gamma)_*\omega_\Gamma$$

where π^∂_Γ is the restriction of π_Γ onto the boundaries of fibers. The second equality holds since ω_Γ is a product of closed forms. So we need to study the boundaries of fibers to prove Theorem 3.4. The proof will be given in §5.

Here we state one more result, which concerns the choices of volume forms.

Proposition 3.5 ([5]). *Suppose g, n and j satisfy (1) or (2) in Theorem 3.4 and $n-j > 2$. Let v_0 and v_1 be two (anti-)symmetric volume forms of S^{n-1} with total integral one, and I_0, I_1 the corresponding integration maps. Then $I_1(\Gamma) - I_0(\Gamma)$ is an exact form for any graph cocycle $\Gamma = \sum a_i\Gamma_i$.*

Proof. Choose any $w' \in \Omega^{n-2}_{DR}(S^{n-1})$ such that $v_1 - v_0 = dw'$, and put $w = (w' + (-1)^n\iota^*_{S^{n-1}}w')/2$. Then $dw = v_1 - v_0$ and $\iota^*_{S^{n-1}}w = (-1)^n w$. Define

$$\tilde{v} := v_0 + d(tw) \in \Omega^{n-1}_{DR}(S^{n-1}\times[0,1]),$$

where t is the coordinate of $[0,1]$. Then $\tilde{v}$ is a (anti-)symmetric closed form on $S^{n-1}\times[0,1]$. Moreover the restriction of $\tilde{v}$ onto $S^{n-1}\times\{\varepsilon\}$, $\varepsilon = 0, 1$, is v_ε.

Assigning to θ-edges e of Γ_i the differential forms

$$\tilde{\theta}_e := (\varphi^\theta_e \times \mathrm{id})^*\tilde{v} \in \Omega^{n-1}_{DR}(C_{\Gamma_i}\times[0,1])$$

instead of θ_e, and integrating their product along the fiber of $\pi_{\Gamma_i}\times\mathrm{id}_{[0,1]}$, we obtain a differential form of $\mathcal{K}_{n,j}\times[0,1]$, which we denote by $\tilde{I}(\Gamma_i)$. Its restriction onto $\mathcal{K}_{n,j}\times\{\varepsilon\}$, $\varepsilon = 0, 1$, is $I_\varepsilon(\Gamma_i)$.

In Lemma 5.25 we will prove that $\tilde{I}(\Gamma) \in \Omega^*_{DR}(\mathcal{K}_{n,j}\times[0,1])$ is a closed form if $n-j > 2$. This completes the proof; if we denote the first projection by p : $\mathcal{K}_{n,j}\times[0,1] \to \mathcal{K}_{n,j}$, then by the generalized Stokes theorem

$$d_{\mathcal{K}_{n,j}}p_*\tilde{I}(\Gamma) = p_*d_{\mathcal{K}_{n,j}\times[0,1]}\tilde{I}(\Gamma) \pm p^\partial_*\tilde{I}(\Gamma) = \pm(\tilde{I}(\Gamma)|_{\mathcal{K}_{n,j}\times\{1\}} - \tilde{I}(\Gamma)|_{\mathcal{K}_{n,j}\times\{0\}})$$

and thus $I_1(\Gamma) - I_0(\Gamma) = \pm dp_* \tilde{I}(\Gamma)$. □

In completely similar way we can prove the following.

Proposition 3.6. *Suppose g, n and j are as in Proposition 3.5. Then the cohomology class $[I(\Gamma)]$ for a graph cocycle Γ does not depend on the choice of $vol_{S^{j-1}}$.*

4. The Haefliger invariant

Here we assume $n-j \geq 3$ is odd, $j \geq 2$ and $2n-3j-3 \geq 0$. Then at present we cannot prove that I is a cochain map, but in this section we describe a $(2n-3j-3)$-cocycle $\mathcal{H} := I(H)+c$ of $\mathcal{K}_{n,j}$ using a graph cocycle $H \in \mathcal{D}_0^{2,0}$ and some 'correction term' c.

The graph cocycle H is shown in Figure 4.1. We call the first graph (with four i-vertices) H_1 and the second H_2; $H = H_1/2 + H_2/6$. By using the rules in Proposition 2.9, we can see that H is indeed a cocycle.



Figure 4.1. Graph cocycle H

Theorem 4.1. *Suppose $n-j \geq 3$ is odd, $j \geq 2$ and $2n-3j-3 \geq 0$. There exists a differential form $c \in \Omega_{DR}^{2n-3j-3}(\mathcal{K}_{n,j})$ such that $\mathcal{H} := I(H)+c$ is closed.*

Though rigorous definition of c and the proof of Theorem 4.1 can be found in §5.9, we give a rough explanation here. Let $\mathcal{I}_j(\mathbb{R}^n)$ be the space of linear injections $\mathbb{R}^j \to \mathbb{R}^n$. Then $c := -p_* D^* \mu$, where $D : \mathcal{K}_{n,j} \times \mathbb{R}^j \to \mathcal{I}_j(\mathbb{R}^n)$ is the derivation map, $p : \mathcal{K}_{n,j} \times \mathbb{R}^j \to \mathcal{K}_{n,j}$ is the first projection and the form $\mu \in \Omega_{DR}^{2n-2j-3}(\mathcal{I}_j(\mathbb{R}^n))$ is given so that $d\mu$ describes the contribution of a boundary stratum of C_{H_2} corresponding to the collision of all the four points. See Definition 5.23 for details.

Below we prove Theorem 1.3, which states that $\mathcal{H}$ gives a non-zero cohomology class and is the Haefliger invariant when $2n-3j-3=0$.

4.1. Additivity.

Here we assume $2n-3j-3=0$, which implies $n=6k$ and $j=4k-1$ for some $k \geq 1$. Then $\mathcal{H} = I(H)+c$ is an isotopy invariant for long $(4k-1)$-knots in $\mathbb{R}^{6k}$. What we will show in this subsection is the 'additivity' of the invariant $\mathcal{H}$.

Proposition 4.2. *The invariant $\mathcal{H}$ is additive under the connect-sum; for any $f_+, f_- \in \mathcal{K}_{6k,4k-1}$, the Kronecker pairing satisfies*

$$\langle \mathcal{H}, f_+ \sharp f_- \rangle = \langle \mathcal{H}, f_+ \rangle + \langle \mathcal{H}, f_- \rangle.$$

Notation. We will show in Lemma 5.26 that the invariant $\mathcal{H}$ is independent of the choice of (anti-)symmetric volume forms. So we may choose (anti-)symmetric volume forms $vol_{S^{N-1}}$, $N = 4k-1$ or $6k$ such that

- their supports are contained in the sufficiently small neighborhoods of the poles $p_{\pm}^{N-1} := (0, \dots, 0, \pm 1) \in S^{N-1}$, and

- they are “$O(N-1)$-invariant,” that is, $A^*vol_{S^{N-1}} = (\det A)vol_{S^{N-1}}$ for any $A \in O(N-1)$ regarded as in $O(N-1) \oplus 1 \subset O(N)$ (in other words $A \in O(N)$ fixes x_N-axis).

For any $f \in \mathcal{K}_{n,j}$, the *support* of f, denoted by $\mathrm{supp}(f) \subset [-1,1]^j$, is defined by

$$\mathrm{supp}(f) := \overline{\{x \in \mathbb{R}^j \mid f(x) \neq (x,0) \in \mathbb{R}^j \times \{0\}^{n-j}\}}.$$

Since we are considering an isotopy invariant, we may suppose $\mathrm{supp}(f_\pm) \subset B^{4k-1}_\pm(\varepsilon)$ and $f_\pm(\mathrm{supp}(f_\pm)) \subset B^{6k}_\pm(\varepsilon/2) \cup (\mathbb{R}^{4k-1} \times \{0\}^{2k+1})$, where

$$B^m_\pm(\varepsilon) := \{(x_1,\dots,x_m) \in \mathbb{R}^m \mid (x_1 \pm 1/2)^2 + x_2^2 + \dots + x_m^2 < \varepsilon^2\},$$

and $\varepsilon > 0$ is a sufficiently small number.

First we compute $\langle \mathcal{H},\, f_+\rangle = \langle I(H_1),\, f_+\rangle/2 + \langle I(H_2),\, f_+\rangle/6 + \langle c,\, f_+\rangle$. The first term $\langle I(H_1),\, f_+\rangle$ is equal to

$$\int_{C_{H_1}(f_+)} \omega_{H_1},$$

where $C_{H_1}(f) \approx C^o_4(\mathbb{R}^{4k-1})$ is the fiber of π_{H_1} over $f \in \mathcal{K}_{6k,4k-1}$. Recall that the integrand is $\theta_{12}\eta_{23}\theta_{34}$ restricted to the fiber. Since we choose $vol_{S^{N-1}}$ with support ‘localized’ around $p^{N-1}_\pm$, we must know for which configurations $x = (x_1,\dots,x_4) \in C_{H_1}(f_+) = C_4(\mathbb{R}^{4k-1})$ all the three vectors $\varphi^\theta_{12}(x)$, $\varphi^\theta_{34}(x)$ and $\varphi^\eta_{23}(x)$ are near $p^{6k-1}_\pm$ or $p^{4k-2}_\pm$; ω_{H_1} does not vanish only on the subspace of such configurations.

Lemma 4.3. *Define* $X_+ := C^o_4(B^{4k-1}_+(\varepsilon)) \subset C_{H_1}(f_+)$. *Then*

$$\langle I(H_1),\, f_+\rangle = \int_{X_+} \omega_{H_1}.$$

Proof. For example, consider $(x_1,\dots,x_4) \in C_{H_1}(f_+) \setminus X_+$ with $x_1 \notin B^{4k-1}_+(\varepsilon)$. Then by our choice of f_+, the vector $\varphi^\theta_{12}(x_1,\dots,x_4)$ cannot be in the support of $vol_{S^{6k-1}}$ (a neighborhood of the pole p^{6k-1}). So the form θ_{12} vanishes on such configurations. In this way we can see that the integrand does not vanish only on X_+, since outside of X_+ the images of φ^θ_{12} and φ^θ_{34} cannot intersect with the supports of $vol_{S^{6k-1}}$ simultaneously. □

Next we compute $\langle I(H_2),\, f_+\rangle$ in similar fashion. This equals

$$\int_{C_{H_2}(f_+)} \omega_{H_2},$$

where $\omega_{H_2} = \theta_{14}\theta_{24}\theta_{34}$. Recall $C_{H_2}(f_+) \subset C^o_3(\mathbb{R}^{4k-1}) \times \mathbb{R}^{6k}$.

Lemma 4.4. *Define a subspace* $Y_+ \subset C_{H_2}(f_+)$ *by*

$$Y_+ := \{(x_1,x_2,x_3;y) \in C_{H_2}(f_+) \mid \text{at least two } x_p \in B^{4k-1}_+(\varepsilon)\}.$$

Then

$$\langle I(H_2),\, f_+\rangle = \int_{Y_+} \omega_{H_2}.$$

Proof. This is because the image of

$$\varphi_{H_2} := \varphi^\theta_{14} \times \varphi^\theta_{24} \times \varphi^\theta_{34} : C_{H_2}(f_+) \setminus Y_+ \longrightarrow (S^{6k-1})^3$$

is of positive codimension, and hence the pullback of the top form of $(S^{6k-1})^3$ must vanish on $C_{H_2}(f_+) \setminus Y_+$. Indeed, outside Y_+ at least two points $f_+(x_p)$, $f_+(x_q)$

are on the standardly embedded $\mathbb{R}^{4k-1}$, hence the image of $C_{H_2} \setminus Y_+$ is the set of $(u_1, u_2, u_3) \in (S^{6k-1})^3$ such that one vector u_p has to be in the linear subspace $(\mathbb{R}^{4k-1} \times \{0\}^{2k+1}) + \mathbb{R}u_q$ of $\mathbb{R}^{6k}$, hence is of codimension $\geq 2k$. $\square$

Now consider the third term $\langle c,\, f_+\rangle$. This is an integration of $D(f_+)^*\mu$ over $\mathbb{R}^{4k-1}$, where $D(f_+) : \mathbb{R}^{4k-1} \to \mathcal{I}_{4k-1}(\mathbb{R}^{6k})$ is given by $x \mapsto (df_+)_x$ (for the definition of μ, see Definition 5.23 and the explanation after Theorem 4.1). Since $D(f_+)$ is constant outside $\mathrm{supp}(f_+)$, we obtain the following.

Lemma 4.5. $\langle c,\, f_+\rangle = \displaystyle\int_{B_+^{4k-1}(\varepsilon)} D(f_+)^*\mu.$

Finally we can compute $\langle \mathcal{H},\, f_+\sharp f_-\rangle$. The long knot $f_+\sharp f_-$ is isotopic to

$$f_+\sharp f_-(x) := \begin{cases} f_\pm(x) & \text{if } x \in B_\pm^{4k-1}(\varepsilon), \\ (x, 0) & \text{if } x \notin B_+^{4k-1}(\varepsilon) \sqcup B_-^{4k-1}(\varepsilon). \end{cases}$$

First, by completely similar arguments to Lemma 4.3, only the configurations $x = (x_1, \dots, x_4)$ such that x_1 and x_2 are in the same $B_+^{4k-1}(\varepsilon)$ or $B_-^{4k-1}(\varepsilon)$, and so is (x_3, x_4), can contribute to $\langle I(H_1),\, f_+\sharp f_-\rangle$. Moreover, if x_2 and x_3 are in the different $B^{4k-1}(\varepsilon)$'s, then the image of the direction map $\varphi_{23}^{\eta}(x)$ cannot be in $\mathrm{supp}(vol_{S^{4k-2}})$. Thus only the configurations with all four x_p's being in the same $B_+^{4k-1}(\varepsilon)$ or $B_-^{4k-1}(\varepsilon)$ can contribute to the pairing, and hence

$$\langle I(H_1),\, f_+\sharp f_-\rangle = \int_{X_+ \sqcup X_-} \omega_{H_1} = \int_{X_+} \omega_{H_1} + \int_{X_-} \omega_{H_1}, \tag{4.1}$$

where $X_- := C_4^o(B_-^{4k-1}(\varepsilon))$.

Next, similarly to Lemma 4.5,

$$\langle c,\, f_+\sharp f_-\rangle = \int_{B_+^{4k-1}(\varepsilon) \sqcup B_-^{4k-1}(\varepsilon)} D(f_+\sharp f_-)^*\mu = \sum_{i=\pm} \int_{B_i^{4k-1}(\varepsilon)} D(f_i)^*\mu. \tag{4.2}$$

As for $I(H_2)$, similar argument to Lemma 4.4 shows

$$\langle I(H_2),\, f_+\sharp f_-\rangle = \int_{Y_+(f_+\sharp f_-)} \omega_{H_2} + \int_{Y_-(f_+\sharp f_-)} \omega_{H_2} + \int_{Y_0(f_+\sharp f_-)} \omega_{H_2}, \tag{4.3}$$

where $Y_-(f_+\sharp f_-) \subset C_{H_2}(f_+\sharp f_-)$ is defined in similar way as Y_+ with B_+'s replaced by B_-'s, and $Y_0(f_+\sharp f_-) \subset C_{H_2}(f_+\sharp f_-)$ consists of $(x_1, x_2, x_3; y)$ with exactly one x_p in each $B_\pm^{4k-1}(\varepsilon)$. By our choice of volume forms, the integrand ω_{H_2} does not vanish only on the subspace consisting of (x_1, x_2, x_3, y) with

1. y is very near '$(0, \dots, 0, \pm\infty)$,' or
2. the first $6k - 1$ factors of $f_+(x_1)$, $f_+(x_2)$, $f_+(x_3)$ and y are close to each other (see Figure 4.2).

On $Y_0(f_+\sharp f_-)$ the condition (2) cannot hold, so we can restrict the integral over $Y_0(f_+\sharp f_-)$ to that over the subspace of $Y_0(f_+\sharp f_-)$ with $|y|$ large. Then since the image of $f_+\sharp f_-$ is very close to that of the trivial knot f_0 (if ε is very small relative to $|y|$),

$$\int_{Y_0(f_+\sharp f_-)} \omega_{H_2} - \int_{C_{H_2}(f_0)} \omega_{H_2} = O(\varepsilon).$$



FIGURE 4.2. Configurations which contribute to $\langle I(H_2),\, f_+\sharp f_-\rangle$

But since $\mathrm{supp}(f_0)=\emptyset$, Lemma 4.4 shows that $\int_{C_{H_2}(f_0)}\omega_{H_2}=0$. Hence

$$\int_{Y_0(f_+\sharp f_-)}\omega_{H_2}=O(\varepsilon). \tag{4.4}$$

$Y_\pm(f_+\sharp f_-)$ is decomposed into two subspaces; in one space, two x_p, x_q are in $B^{4k-1}_\pm(\varepsilon)$ and remaining x_r is in $B^{4k-1}_\mp(\varepsilon)$, and in the other space x_r is not in $B^{4k-1}_\mp(\varepsilon)$. In the former space, the condition (2) above cannot hold, so $|y|$ has to be large. So replacing $f_+\sharp f_-$ with $f_\pm$ changes the integral by $O(\varepsilon)$;

$$\int_{Y_\pm(f_+\sharp f_-)}\omega_{H_2}=\int_{Y_\pm(f_\pm)}\omega_{H_2}+O(\varepsilon). \tag{4.5}$$

Summing up (4.1), (4.2) and (4.3) and substituting (4.4) and (4.5), we have

$$\begin{aligned}\langle I(H),\, f_+\sharp f_-\rangle &= \sum_{i=\pm}\left(\frac{1}{2}\int_{X_i}\omega_{H_1}+\frac{1}{6}\int_{Y_i(f_i)}\omega_{H_2}+\int_{B_i^{4k-1}(\varepsilon)}D(f_i)^*\mu\right)+O(\varepsilon)\\ &= \langle I(H),\, f_+\rangle+\langle I(H),\, f_-\rangle+O(\varepsilon).\end{aligned}$$

This completes the proof of Proposition 4.2, since $\langle\mathcal{H},\, f_+\sharp f_-\rangle$ is independent of ε.

Remark 4.6. By similar argument, the order two invariant [14, 20] for $\mathcal{K}_{m+2,m}$ ($m\geq 3$ is odd) is proved to be additive under the connect-sum.

4.2. **Evaluation.**

Here we prove the last part of Theorem 1.3.

We know by [9] that $\pi_0(\mathcal{K}_{6k,4k-1})$ forms a group under the connect-sum and is isomorphic to $\mathbb{Z}$. Our goal here is the following.

Theorem 4.7. *When $n=6k$ and $j=4k-1$ for some $k\geq 1$, then $\langle\mathcal{H},\, S\rangle$ is equal to ± 1 for the generator S of $\pi_0(\mathcal{K}_{6k,4k-1})$.*

Since both the Haefliger invariant and our $\mathcal{H}$ are additive under the connect-sum (see [8] and Proposition 4.2), Theorem 4.7 says that $\mathcal{H}$ is the Haefliger invariant.

We will use the generator of $\pi_0(\mathcal{K}_{6k,4k-1})$ given by Budney [4] and Roseman-Takase [13].

4.2.1. *Deform-spinning.* Given an N-fold based loop $\gamma\in\Omega^N\mathcal{K}_{n,j}$ (for any n,j) represented by a smooth map $\gamma:\mathbb{R}^N\to\mathcal{K}_{n,j}$ with $\gamma(x)\equiv\iota$ (the trivial long j-knot) for any $x\notin[-1,1]^j$, we have $S_\gamma\in\mathcal{K}_{N+n,N+j}$ defined by

$$S_\gamma(x,t):=(x,\gamma(x)(t))\in\mathbb{R}^N\times\mathbb{R}^n,\quad x\in\mathbb{R}^N,\ t\in\mathbb{R}^j.$$

This is a special case of 'deform-spinning' construction [12]. Putting $N=1$, we obtain the *graphing map* $\mathrm{gr}:\Omega\mathcal{K}_{n,j}\to\mathcal{K}_{n+1,j+1}$ given in [4]. It has been shown in

[4] that all the following maps

$$\pi_{4k-2}(\mathcal{K}_{2k+2,1}) \xrightarrow{\mathrm{gr}} \pi_{4k-3}(\mathcal{K}_{2k+3,2}) \xrightarrow{\mathrm{gr}} \dots \xrightarrow{\mathrm{gr}} \pi_0(\mathcal{K}_{6k,4k-1}) \cong \mathbb{Z}$$

are isomorphisms, so $\pi_0(\mathcal{K}_{6k,4k-1}) \cong \mathbb{Z}$ is generated by $S_\psi := \mathrm{gr}^{4k-2}(\psi)$;

$$S_\psi(z) := (x, \psi(x)(t)) \in \mathbb{R}^{4k-2} \times \mathbb{R}^{2k+2}, \quad z = (x,t) \in \mathbb{R}^{4k-2} \times \mathbb{R}^1, \tag{4.6}$$

where $\psi : \mathbb{R}^{4k-2} \to \mathcal{K}_{2k+2,1}$ is a map which gives the generator of $\pi_{4k-2}(\mathcal{K}_{2k+2,1})$. In fact $\pi_{4k-2}(\mathcal{K}_{2k+2,1})$ is the first non-vanishing homotopy group of $\mathcal{K}_{2k+2,1}$ (for example see [4, 18]), and the image of its generator ψ via Hurewicz isomorphism is given as follows (probably the dual cocycle to ψ first appeared in [19]; see also [4, 5, 11, 15]).

Consider a 'long immersion' $f : \mathbb{R}^1 \to \mathbb{R}^{2k+2}$ which has only two transversal doublepoints $z_i = f(\xi_i) = f(\xi_{i+2})$, $i = 1, 2$, $\xi_1 < \xi_2 < \xi_3 < \xi_4$ (see Figure 4.3). Consider the unit sphere S_i^{2k-1} in the complementary $2k$-space to $\mathbb{R}f'(\xi_i) \oplus \mathbb{R}f'(\xi_{i+2})$ in $T_{z_i}\mathbb{R}^{2k+2}$. At each doublepoint z_i, we can 'lift' the segments $f(\xi_{i+2} - \varepsilon, \xi_{i+2} + \varepsilon)$ to the directions of complementary S_i^{2k-1} to obtain non-singular embeddings (see Figure 4.4). More precisely, for $(w_1, w_2) \in (S^{2k-1})^2$, define $\psi(w_1, w_2) \in \mathcal{K}_{2k+2,1}$ by

$$\psi(w_1, w_2)(t) := \begin{cases} f(t) + \left(\delta \exp\left[\frac{1}{(t-\xi_{i+2})^2 - \varepsilon^2}\right]\right) w_i & |t - \xi_{i+2}| < \varepsilon,\ i = 1, 2 \\ f(t) & \text{otherwise} \end{cases}$$

where δ and ε are positive small numbers. Thus a map $\psi = \psi_\delta : (S^{2k-1})^2 \to \mathcal{K}_{2k+2,1}$ is defined, and it is known that ψ induces an isomorphism $\psi_* : H_{4k-2}((S^{2k-1})^2) \xrightarrow{\cong} H_{4k-2}(\mathcal{K}_{2k+2,1})$.



FIGURE 4.3. Immersion f in $\mathbb{R}^{2k+2}$



FIGURE 4.4. Definition of ψ

The following is proved by inspection (see Figure 4.4).

Lemma 4.8. *The union*

$$W_3 := \bigcup_{w_1 \in S^{2k-1}} \psi(w_1, w_2)(\xi_3 - \varepsilon, \xi_3 + \varepsilon) \subset \mathbb{R}^{2k+2}$$

is homeomorphic to S^{2k}. *Consider the map*

$$\varphi : W_3 \times I_1 \longrightarrow S^{2k+1}, \quad (x, y) \longmapsto \frac{y - x}{|y - x|},$$

where $I_1 := f(\xi_1 - \varepsilon, \xi_1 + \varepsilon)$. *The linking number of* W_3 *with* I_1 *is one; more precisely, the limit* $\delta \to 0$ *of the integration of* $\varphi^* vol_{S^{2k+1}}$ *over* $W_3 \times I_1$ *is* ± 1. *Similar statement holds for*

$$W_4 := \bigcup_{w_2 \in S^{2k-1}} \psi(w_1, w_2)(\xi_4 - \varepsilon, \xi_4 + \varepsilon) \quad \text{and} \quad I_2 := f(\xi_2 - \varepsilon, \xi_2 + \varepsilon).$$

Fixing the natural projection $\pi : [-1/2, 1/2]^{2k-1} \to S^{2k-1}$ such that

- $q^{2k-1}_{\pm} := \pi^{-1}(p^{2k-1}_{\pm}) \in (-1/2, 1/2)^{2k-1}$ (recall $p^{2k-1}_{\pm} := (0, \dots, 0, \pm 1) \in S^{2k-1}$),
- $\pi(\partial[-1/2, 1/2]^{2k-1})$ consists of a single point $u_0 := (1, 0, \dots, 0)$,

we can think of ψ as $\psi : [-1/2, 1/2]^{4k-2} \to \mathcal{K}_{2k+2,1}$. We can extend ψ to whole $\mathbb{R}^{4k-2}$ so that $\psi(x) \equiv \iota$ if $x \notin [-1, 1]^{4k-2}$ (see also §4.2.2), since $\mathcal{K}_{2k+2,1}$ is $(4k-3)$-connected as mentioned above. Thus we have $S_\psi \in \mathcal{K}_{6k,4k-1}$ defined as (4.6).

Theorem 4.9 ([4, 13])**.** *If* ψ *is as above, then* S_ψ *generates* $\pi_0(\mathcal{K}_{6k,4k-1})$.

Remark 4.10. In [13] a generator of $\pi_0(\mathrm{Emb}\,(S^{4k-1}, S^{6k}))$ was defined by the deform-spinning $\psi : (S^{2k-1})^2 \to \mathcal{K}_{2k+2,1}$ along the torus. But it is not difficult to see that such a spinning also gives an element of $\mathcal{K}_{6k,4k-1}$ which is isotopic to our S_ψ given by the graphing map [4].

The above construction can be done for the generator of $\pi_{2d-2}(\mathcal{K}_{d+2,1})$ for any $d > 1$ to obtain $S_\psi \in \mathcal{K}_{3d,2d-1}$. Since we assumed that $n - j = d + 1$ is odd, we put $d = 2k$ here (otherwise there exist orientation reversing automorphisms of graphs H_i, and hence they vanish). When d is odd, any generator of $\pi_0(\mathcal{K}_{3d,2d-1}) \cong \mathbb{Z}/2$ would not be detected by any de Rham cohomology classes.

4.2.2. *A suitable choice of* $\psi : \mathbb{R}^{4k-2} \to \mathcal{K}_{2k+2,1}$. Below we will compute

$$\langle \mathcal{H}, S_\psi \rangle = \frac{1}{2} \int_{C_{H_1}(S_\psi)} \omega_{H_1} + \frac{1}{6} \int_{C_{H_2}(S_\psi)} \omega_{H_2} + \int_{\mathbb{R}^{4k-1}} D(S_\psi)^* \mu, \tag{4.7}$$

where $D(S_\psi) : \mathbb{R}^{4k-1} \to \mathcal{I}_{4k-1}(\mathbb{R}^{6k})$ is defined by $D(S_\psi)(x) := (dS_\psi)_x$, and μ is defined in the sentence after the proof of Lemma 5.22. To simplify the computation of (4.7), we need to choose a favorable extension $\psi : \mathbb{R}^{4k-2} \to \mathcal{K}_{2k+2,1}$.

First, we take the immersion f as in Figure 4.3; almost all the image of $\psi(x)$ is contained in the 2-plane $\mathbb{R}^2 \times \{0\}^{2k} \subset \mathbb{R}^{2k+2}$, except for the neighborhoods of z_1, z_2 and another crossing c which corresponds to $f(\zeta_1)$ and $f(\zeta_2)$, $\xi_2 < \zeta_1 < \xi_3$, $\xi_4 < \zeta_2 < 1$. We suppose that the arc $\psi(x)(\zeta_1 - \varepsilon, \zeta_1 + \varepsilon)$ is in the 2-plane, $\psi(x)(\zeta_2 - \varepsilon, \zeta_2 + \varepsilon)$ is in the 3-plane $\mathbb{R}^3 \times \{0\}^{2k-1} \subset \mathbb{R}^{2k+2}$, and that the resolutions of segments $\psi(x)(\xi_{i+2} - \varepsilon, \xi_{i+2} + \varepsilon)$ occur in $\{0\}^2 \times \mathbb{R}^{2k}$.

Next we suppose that the image of $\psi(x)$ is 'almost' in $\mathbb{R}^2$;

$$\psi(x)(\mathbb{R}^1) \subset \mathbb{R}^2 \times (-\delta, \delta)^{2k} \quad \text{for any } x \in \mathbb{R}^{4k-2}. \tag{4.8}$$

Lastly we suppose ψ is 'symmetric' in the following sense; define

$$\begin{aligned} X &:= \{(s,t,u) \in \mathbb{R}^{2k-1} \times \mathbb{R}^{2k-1} \times \mathbb{R}^1 \mid s \notin [-1/2,1/2]^{2k-1}, t \in [-1/2,1/2]^{2k-1}\}, \\ Y &:= \{(s,t,u) \in \mathbb{R}^{2k-1} \times \mathbb{R}^{2k-1} \times \mathbb{R}^1 \mid s \in [-1/2,1/2]^{2k-1}, t \notin [-1/2,1/2]^{2k-1}\}, \\ Z &:= \{(s,t,u) \in \mathbb{R}^{2k-1} \times \mathbb{R}^{2k-1} \times \mathbb{R}^1 \mid s,t \notin [-1/2,1/2]^{2k-1}\} \end{aligned}$$

and X', Y', Z' as the images of X, Y, Z respectively via the first $4k-2$ projection $pr_{4k-2} : \mathbb{R}^{4k-1} \to \mathbb{R}^{4k-2}$ (see Figure 4.5). Define $\psi : X' \to \mathcal{K}_{2k+2,1}$ by

$$\psi(s,t) = h_{|s|/|s'|}(\pi(t))$$

where $h_\alpha : S^{2k-1} \to \mathcal{K}_{2k+2,1}$ ($\alpha \geq 1$) is a one parameter family with $h_1 = \psi(u_0, -)$ and $h_\alpha \equiv \iota$ (the trivial long knot) for $\alpha \geq 2$, and $s' \in \partial[-1/2,1/2]^{2k-1}$ is the unique element with $s = \alpha s'$ for some $\alpha \geq 1$. Using similar one parameter family connecting $\psi(-, u_0)$ (resp. $\psi(u_0,u_0)$) and ι, we can define ψ on Y' (resp. on Z') (see Example 4.11 and Figure 4.5). We can modify ψ so that it is smooth. The following properties of ψ will be important below;

$$\psi(s,t) = \psi(-s,t) \quad \text{if } (s,t) \in X' \cup Z', \tag{4.9}$$

$$\psi(s,t) = \psi(s,-t) \quad \text{if } (s,t) \in Y' \cup Z'. \tag{4.10}$$

Example 4.11. Consider the case $k = 1$ (see Figure 4.5). $X' \to \mathcal{K}_{4,1}$ is given as



FIGURE 4.5. Extension $\psi : \mathbb{R}^2 \to \mathcal{K}_{4,1}$

a null homotopy $\psi(s,-) : S^1 \to \mathcal{K}_{4,1}$ ($|s| \geq 1/2$) from $\psi(\{1/2\} \times S^1)$ to ι. Since $\psi(\{1/2\} \times [-1/2,1/2])$ and $\psi(\{-1/2\} \times [-1/2,1/2])$ are the same cycle of $\mathcal{K}_{4,1}$, two homotopies corresponding to $X' \cap \{s \geq 1/2\}$ and $X' \cap \{s \leq -1/2\}$ can be taken so that $\psi(s,t) = \psi(-s,t)$. This is (4.9). Similarly $Y' \to \mathcal{K}_{4,1}$ is a homotopy from $\psi(S^1 \times \{1/2\})$ to ι, and $\psi(s,t) = \psi(s,-t)$ on Y'.

Eight thick segments $\{\pm 1/2\} \times \{|t| \geq 1/2\}$, $\{|s| \geq 1/2\} \times \{\pm 1/2\}$ in Figure 4.5 represent the same homotopy from $\psi(1/2,1/2)$ to ι. Thus we can define ψ by

$$\psi(r\cos\beta + 1/2, r\sin\beta + 1/2) := \psi(r + 1/2, 1/2) \quad (0 \leq \beta \leq \pi/4)$$

on $Z' \cap \{s,t \geq 1/2\}$, and similarly on other components. Then it is easy to see $\psi(s,t) = \psi(-s,t) = \psi(s,-t) = \psi(-s,-t)$ on Z'.

Using such an extension ψ, we will compute (4.7).

4.2.3. *The first term* $\langle I(H_1),\, S_\psi\rangle$. First let us study on which configurations $x = (x_1,\dots,x_4)\in C_{H_1}(S_\psi)$ the integrand ω_{H_1} does not vanish, as was done in §4.1.

Let us write $x_p := (s^{(p)}, t^{(p)}, u^{(p)}) \in (\mathbb{R}^{2k-1})^2\times\mathbb{R}^1$ $(p=1,\dots,4)$. Then by (4.6),

$$\begin{aligned} &S_\psi(x_2)-S_\psi(x_1) = \\ &\quad (s^{(2)}-s^{(1)}, t^{(2)}-t^{(1)}, \psi(s^{(2)},t^{(2)})(u^{(2)})-\psi(s^{(1)},t^{(1)})(u^{(1)})) \\ &\qquad\qquad\qquad\qquad\in (\mathbb{R}^{2k-1})^2\times\mathbb{R}^{2k+2}. \end{aligned} \tag{4.11}$$

By (4.8), the length of the last $2k$ factors of (4.11) is at most $2\sqrt{2k}\,\delta$. So $\varphi^\theta_{12}(x_1,x_2) = (S_\psi(x_2)-S_\psi(x_1))/|S_\psi(x_2)-S_\psi(x_1)|$ is near $p^{6k-1}_\pm$ only if at least the first $4k-2$ factors of (4.11) are nearly zero, that is, $(s^{(1)},t^{(1)})$ is near $(s^{(2)},t^{(2)})$. Similarly $(s^{(3)},t^{(3)})$ must be near $(s^{(4)},t^{(4)})$. Moreover $(s^{(2)},t^{(2)})$ must be near $(s^{(3)},t^{(3)})$, since $w=(x_3-x_2)/|x_3-x_2|$ is near $p^{4k-2}_\pm$. Thus we need to consider only $x=(x_1,\dots,x_4)$ with all $(s^{(p)},t^{(p)})$ $(p=1,\dots,4)$ close to each other. Notice that they must become closer and closer if we choose $vol_{S^{N-1}}$ with smaller support.

Let $L := (-1/2-a, 1/2+a)^{4k-2}\times\mathbb{R}^1$ ($a>0$ is a fixed small number) be a small neighborhood of $[-1/2,1/2]^{4k-2}\times\mathbb{R}^1$, and consider $x\in C_4(\mathbb{R}^{4k-1}\setminus L)$.

Lemma 4.12. *If we choose ψ as in §4.2.2 and $vol_{S^{N-1}}$ with sufficiently small support, then the integration of ω_{H_1} over $C_4(\mathbb{R}^{4k-1}\setminus L)$ vanishes.*

Proof. If $\mathrm{supp}(vol_{S^{N-1}})$ is sufficiently small relative to $\delta>0$, then $(x_1,\dots,x_4)\in C_4(\mathbb{R}^{4k-1}\setminus L)$ must be such that all $(s^{(p)},t^{(p)})$ $(p=1,\dots,4)$ are close to each other so that at most one of $\{x_1,\dots,x_4\}\cap(X\setminus L)$ and $\{x_1,\dots,x_4\}\cap(Y\setminus L)$ is not empty.

Consider the integration of ω_{H_1} over $\{x\in C_4(\mathbb{R}^{4k-1}\setminus L)\,|\,x\cap X\neq\emptyset\}$. Notice that in this case $x\in C_4(X\cup Z)$. Define an involution F_1 of this subspace by

$$F_1(x_1,\dots,x_4) := (i_1x_1,\dots,i_1x_4),$$

where $i_1:\mathbb{R}^{4k-1}\to\mathbb{R}^{4k-1}$ is given by $i_1(s,t,u) := (-s,t,u)$ $(s,t\in\mathbb{R}^{2k-1},\ u\in\mathbb{R}^1)$. The map F_1 preserves the orientation, but $F_1^*\omega_{H_1} = -\omega_{H_1}$ because

- $\varphi^\theta_{12}\circ F_1 = i_1\circ\varphi^\theta_{12}$ by (4.9) and (4.11) (here we abbreviate $i_1\times\mathrm{id}_{\mathbb{R}^{2k+1}}:\mathbb{R}^{6k}\to\mathbb{R}^{6k}$ to i_1), and hence $F_1^*\theta_{12} = (-1)^{2k-1}\theta_{12} = -\theta_{12}$ (since we have assumed $i_1^*vol_{S^{6k-1}} = (-1)^{2k-1}vol_{S^{6k-1}}$; see the remark after Proposition 4.2),
- similarly $F_1^*\theta_{34} = -\theta_{34}$, and
- $\varphi^\eta_{23}\circ F_1 = i_1\circ\varphi^\eta_{23}$ and hence $F_1^*\eta_{23} = -\eta_{23}$.

Hence the integration of ω_{H_1} over $\{x\in C_4(\mathbb{R}^{4k-1}\setminus L)\,|\,x\cap X\neq\emptyset\}$ must vanish. Similarly, we can show the vanishing of the integration of ω_{H_1} over $\{x\cap Y\neq\emptyset\}$ by considering an involution F_2 given by $F_2(x_1,\dots,x_4) := (i_2x_1,\dots,i_2x_4)$, where $i_2:\mathbb{R}^{4k-1}\to\mathbb{R}^{4k-1}$ is given by $i_2(s,t,u) := (s,-t,u)$. The vanishing of the integration of ω_{H_1} over $\{x\cap X = x\cap Y=\emptyset\}$ is proved in similar way; in this case $x\in C_4(Z\setminus L)$, and hence either F_1 or F_2 can work. □

By Lemma 4.12, only $x=(x_1,\dots,x_4)$ with at least one x_p in L (and other x_q's near L) contributes to $\langle I(H_1),\, S_\psi\rangle$.

Consider the tubular neighborhood $N=\mathbb{R}^{4k-2}\times[-1/2,\,1/2]$ of $\mathbb{R}^{4k-2}$ in $\mathbb{R}^{4k-1}$. Define four 'fat planes' by

$$N_i := \mathbb{R}^{4k-2}\times(\xi_i-\varepsilon,\,\xi_i+\varepsilon)\subset N\subset\mathbb{R}^{4k-1},\quad 1\le i\le 4$$

(ξ_i and ε have appeared in the definition of ψ). Recall that we write $q_{\pm}^{2k-1} := \pi^{-1}(p_{\pm}^{2k-1})$ (we choose π so that $q_{\pm}^{2k-1} \in (-1/2, 1/2)^{2k-1}$).

Lemma 4.13. *Let $x_1, x_2 \in \mathbb{R}^{4k-1}$ be two distinct points with at least one $x_p \in L$. Then the direction $\varphi_{12}^{\theta}(x_1, x_2)$ is near $p_{\pm}^{6k-1}$ only if*

- *$(s^{(1)}, t^{(1)})$ is near $(s^{(2)}, t^{(2)})$,*
- *$x_1 \in N_i$ and $x_2 \in N_{i+2}$ for some $i = 1, \dots, 4$ (here the suffixes are understood modulo 4), and*
- *$s^{(2)}$ (resp. $t^{(2)}$, $s^{(1)}$, $t^{(1)}$) is near $q_{\pm}$ when $(x_1, x_2) \in N_1 \times N_3$ (resp. $N_2 \times N_4$, $N_3 \times N_1$, $N_4 \times N_2$).*

Proof. We have already seen that $(s^{(1)}, t^{(1)})$ must be near $(s^{(2)}, t^{(2)})$. But it is not enough; the direction determined by the last $2k+2$ factors $\psi(s^{(2)}, t^{(2)})(u^{(2)}) - \psi(s^{(1)}, t^{(1)})(u^{(1)})$ of (4.11) must be near $p_{\pm}^{2k+1}$, and it is the case only if the two points $\psi(s^{(p)}, t^{(p)})(u^{(p)})$ ($p = 1, 2$) are around a self-intersection z_i of f (see Figure 4.4) which is resolved in a direction near $p_{\pm}^{2k+1}$. Such a situation is realized, for example, if

- $u^{(1)} \in (\xi_1 - \varepsilon, \xi_1 + \varepsilon)$ and $u^{(2)} \in (\xi_3 - \varepsilon, \xi_3 + \varepsilon)$, and
- $\pi(s^{(2)})$ is near $p_{\pm}^{2k+1}$ (recall $\pi : [-1/2, 1/2]^{2k-1} \twoheadrightarrow S^{2k-1}$).

This is the case of $(x_1, x_2) \in N_1 \times N_3$ in the Lemma. $\square$

Thus it suffices to consider $x \in C_{H_1}(S_{\psi})$ such that both (x_1, x_2) and (x_3, x_4) satisfy the condition of Lemma 4.13, and all the $(s^{(p)}, t^{(p)}) \in \mathbb{R}^{4k-2}$ are close to each other. We divide such configurations into two types.

Type I. All the four points $S_{\psi}(x_p)$ ($1 \le p \le 4$) are near the resolution of a single z_i ($i = 1$ or 2).

Let us write $N_{p,q,r,s} := N_p \times N_q \times N_r \times N_s \subset C_4(\mathbb{R}^{4k-1})$ ($p, q, r, s = 1, 2, 3, 4$). Type I configuration $x = (x_1, \dots, x_4)$ is in $N_{i,i+2,i,i+2}$ or $N_{i,i+2,i+2,i}$ for some i (modulo 4). Figure 4.6 shows an example of $x \in N_{1,3,1,3}$ such that $\varphi_{H_1}(x) = (v_1, v_2, w)$ ($\varphi_{H_1} := \varphi_{12}^{\theta} \times \varphi_{34}^{\theta} \times \varphi_{23}^{\eta}$) is in the support of the volume forms.



FIGURE 4.6. $x = \varphi_{H_1}^{-1}(v_1, v_2, w) \in N_{1,3,1,3}$

Type II. Two points $S_{\psi}(x_1)$ and $S_{\psi}(x_2)$ are near the resolution of z_i of f, and $S_{\psi}(x_3)$ and $S_{\psi}(x_4)$ are near the resolution of z_{i+1}, $i = 1$ or 2 (here $z_3 := z_1$).

Such a configuration is in $N_{i,i+2,i+1,i+3}$ or $N_{i,i+2,i+3,i+1}$ for some i. Type II inverse image of $(v_1, v_2, w) \in (S^{6k-1})^2 \times S^{4k-2}$ via φ_{H_1} looks like Figure 4.7, because of the following.

Lemma 4.14. *Let $v \in S^{6k-1}$ near $p_{\pm}^{6k-1}$ be given. If the pair $(x_1, x_2) \in N_1 \times N_3$ satisfies $\varphi_{12}^{\theta}(x_1, x_2) = v$, then $s^{(2)}$ and $u^{(2)}$ are uniquely determined by v. If moreover we give any $t^{(2)}$ to fix x_2, then accordingly x_1 is determined.*

Similarly, if $(x_1, x_2) \in N_2 \times N_4$ satisfies $\varphi_{12}^{\theta}(x_1, x_2) = v$, then (x_1, x_2) is uniquely determined according to given $s^{(2)}$.

Proof. Consider the case $(x_1, x_2) \in N_1 \times N_3$. The last $2k+2$ factors of $\varphi_{12}^{\theta}(x_1, x_2)$ are determined by $\psi(s^{(2)}, t^{(2)})(u^{(2)}) - \psi(s^{(1)}, t^{(1)})(u^{(1)})$ as we see in (4.11). Comparing them with those of v, we can see that $s^{(2)}$, $u^{(1)}$ and $u^{(2)}$ are uniquely determined so that $s^{(2)}$ is near q_+ (see also Lemma 4.8).

Thus we can fix x_2 if we give any $t^{(2)}$. Then x_1 is also uniquely determined, since $u^{(1)}$ is already determined as above, and $(s^{(1)}, t^{(1)})$ is determined by comparing the first $4k-1$ factors of v with those of $\varphi_{12}^{\theta}(x_1, x_2)$. $\square$

Figure 4.7 shows an example of $x = (x_1, \dots, x_4) \in N_{1,3,4,2}$. In this case $x_1, \dots, x_4$ are uniquely determined by $v_1 = \varphi_{12}^{\theta}(x)$ and $v_2 = \varphi_{34}^{\theta}(x)$ up to $t^{(2)}$ and $s^{(4)}$ by Lemma 4.14, and $t^{(2)}$ and $s^{(4)}$ are determined by $w = (x_3 - x_2)/|x_3 - x_2|$.



FIGURE 4.7. $x = \varphi_{H_1}^{-1}(v_1, v_2, w) \in N_{1,3,4,2}$

Firstly we compute the contribution of Type II configurations. Choose neighborhoods $U_{\pm}^{N-1}$ of $p_{\pm}^{N-1}$ ($N = 6k$ or $4k-1$) so that the support of $vol_{S^{N-1}}$ is contained in $U_{+}^{N-1} \sqcup U_{-}^{N-1}$. Also choose neighborhoods $V_{\pm}^{2k-1}$ of $q_{\pm}$ so that all the four points $(x_1, \dots, x_4)$ of Type II configurations are mapped into the same $V_l^{2k-1} \times V_m^{2k-1}$ ($l, m = \pm$) by $pr_{4k-2} : \mathbb{R}^{4k-1} \to \mathbb{R}^{4k-2}$. Define

$$N_i^{l,m} := V_l^{2k-1} \times V_m^{2k-1} \times (\xi_i - \varepsilon,\, \xi_i + \varepsilon) \subset N_i, \quad l, m = \pm.$$

Define $N_{p,q,r,s}^{l,m} := N_p^{l,m} \times N_q^{l,m} \times N_r^{l,m} \times N_s^{l,m}$ ($l, m = \pm$, $\{p, q, r, s\} = \{1, 2, 3, 4\}$). Type II configurations are in $N_{i,i+2,i+1,i+3}^{l,m}$ or $N_{i,i+2,i+3,i+1}^{l,m}$ for some $i = 1, \dots, 4$ and $l, m = \pm$. There are $4 \times 4 \times 2 = 32$ such components. Via the direction map φ_{H_1}, each component is mapped to some $U_l^{6k-1} \times U_{l'}^{6k-1} \times U_{l''}^{4k-2}$ homeomorphically. Hence each component contributes to $\langle I(H_1),\, S_{\psi} \rangle$ by $\pm(1/2)^3 = \pm 1/8$.

The signs are given as follows.

Lemma 4.15. (1) *All the signs for $N_{p,q,r,s}^{l,m}$, $l, m = \pm$, are same for fixed p, q, r, s.*

(2) *If the orientations are chosen so that the sign for $N_{1,3,2,4}^{l,m}$ is $+1$, then the signs for $N_{2,4,1,3}^{l,m}$ and $N_{3,1,4,2}^{l,m}$ are -1, and those for other five components $N_{2,4,3,1}^{l,m}$, $N_{3,1,4,2}^{l,m}$, $N_{4,2,1,3}^{l,m}$, $N_{1,3,4,2}^{l,m}$ and $N_{4,2,3,1}^{l,m}$ are $+1$.*

Proof. (1) is because the difference between the integrations of ω_{H_1} over $N^{l,m}_{p,q,r,s}$ and $N^{-l,m}_{p,q,r,s}$ comes from the antipodal map of S^{2k-1}, which preserves the orientation.

For (2), first we show that the signs for $N^{l,m}_{1,3,2,4}$ and $N^{l,m}_{3,1,4,2}$ are different. This follows from the diagram which is commutative up to isotopy

$$\begin{tikzcd}
N^{l,m}_{1,3,2,4} \arrow[r, "\varphi_{H_1}"] \arrow[d, "G"'] & U^{6k-1}_{l} \times U^{6k-1}_{m} \times U^{4k-2}_{-} \arrow[d, "\iota_{S^{6k-1}} \times \iota_{S^{6k-1}} \times g"] \\
N^{l,m}_{2,4,1,3} \arrow[r, "\varphi_{H_1}"] & U^{6k-1}_{l} \times U^{6k-1}_{m} \times U^{4k-2}_{-}
\end{tikzcd}$$

where

$$G(x_1, \dots, x_4) := (t^{(1)}, s^{(1)}, u^{(1)} - \xi_1 + \xi_2\, ;\, t^{(2)}, s^{(2)}, u^{(2)} - \xi_3 + \xi_4\, ;\\ t^{(3)}, s^{(3)}, u^{(3)} - \xi_3 + \xi_1\, ;\, t^{(4)}, s^{(4)}, u^{(4)} - \xi_4 + \xi_3),$$

and g is induced from $(s, t, u) \mapsto (s, t, -u) \in (\mathbb{R}^{2k-1})^2 \times \mathbb{R}^1$. G preserves the orientation but $\iota_{S^{6k-1}} \times g \times \iota_{S^{6k-1}}$ does not, so the signs for $N^{l,m}_{1,3,2,4}$ and $N^{l,m}_{3,1,4,2}$ are different. The signs for $N^{l,m}_{2,4,1,3}$ and $N^{l,m}_{3,1,4,2}$ are same, since their difference comes from an automorphism of H_1, which always preserves the orientation.

Next we show that all the signs for other five components are same as that for $N^{l,m}_{1,3,2,4}$.

(i) $N^{l,m}_{1,3,2,4}$ and $N^{l,m}_{4,2,3,1}$; the signs for them are same, since the difference arises from an automorphism of H_1.

(ii) $N^{l,m}_{1,3,2,4}$ and $N^{l,m}_{2,4,3,1}$; their signs are same because of the following commutative diagram (up to isotopy);

$$\begin{tikzcd}
N^{l,m}_{1,3,2,4} \arrow[r, "\varphi_{H_1}"] \arrow[d, "G'"'] & U^{6k-1}_{l} \times U^{6k-1}_{m} \times U^{4k-2}_{-} \arrow[d, "i_1 \times \mathrm{id} \times g"] \\
N^{l,m}_{2,4,3,1} \arrow[r, "\varphi_{H_1}"] & U^{6k-1}_{l} \times U^{6k-1}_{m} \times U^{4k-2}_{+}
\end{tikzcd}$$

where

$$G'(x_1, \dots, x_4) := (s^{(2)}, t^{(1)}, u^{(2)}\, ;\, s^{(1)}, t^{(2)}, u^{(1)};\\ s^{(3)} - s^{(2)} + s^{(1)}, t^{(3)}, u^{(3)}\, ;\, s^{(4)} - s^{(2)} + s^{(1)}, t^{(4)}, u^{(4)}),$$

and i_1 was given in Lemma 4.12. The vertical arrows preserve the orientations, so the signs are same. Similar diagrams shows that the signs for $N^{l,m}_{1,3,2,4}$ and $N^{l,m}_{3,1,2,4}$ are same.

(iii) $N^{l,m}_{4,2,1,3}$ and $N^{l,m}_{1,3,4,2}$; their signs are same as those of $N^{l,m}_{3,1,2,4}$ and $N^{l,m}_{1,3,2,4}$ respectively, since their differences come from automorphisms of H_1. □

By Lemma 4.15, 24 components contribute to $\langle I(H_1), S_\psi \rangle$ by ± 1 and other 8 components by ∓ 1. Thus their contribution is $\pm(1/8) \times (24 - 8) = \pm 2$.

Next consider Type I configurations.

Lemma 4.16. *The contribution of Type I configurations is zero.*

Proof. Consider the case when all the four points $S_\psi(x_p)$ are near z_1. Then $x \in C_4((N_1 \sqcup N_3) \cap L)$. But on this space, the direction map φ_{H_1} is invariant under the translation

$$(x_1, \dots, x_4) \mapsto (\tau_v x_1, \dots, \tau_v x_4), \quad v \in \mathbb{R}^{2k-1}$$

where $\tau_v(s,t,u) := (s, t+v, u) \in \mathbb{R}^{2k-1} \times \mathbb{R}^{2k-1} \times \mathbb{R}^1$. This is because of (4.11) and $\psi(s,t)(u) = \psi(s, t+v)(u)$ if $(s,t,u) \in (N_1 \sqcup N_3) \cap L$. Hence the image of φ_{H_1} must be of positive codimension and the integrand $\omega_{H_1} = \varphi_{H_1}^*(vol_{S^{6k-1}} \times vol_{S^{6k-1}} \times vol_{S^{4k-2}})$ must vanish on $C_4((N_1 \sqcup N_3) \cap L)$.

In the case when four points $S_\psi(x_p)$ are near z_2, a similar translation $\tau'_v(s,t,u) := (s+v, t, u)$ would prove the vanishing of ω_{H_1} on $C_4((N_2 \sqcup N_4) \cap L)$. $\square$

Thus the first term of (4.7) is equal to $\pm 2 \times (1/2) = \pm 1$.

4.2.4. *Remaining terms.* To complete the proof of Theorem 4.7, we will prove that the second and the third terms of (4.7) do not contribute to $\langle \mathcal{H}, S_{\psi_\delta} \rangle$.

Contribution of H_2. Now we compute the second term of (4.7). Recall $C_{H_2}(S_\psi) \subset C_3^o(\mathbb{R}^{4k-1}) \times \mathbb{R}^{6k}$.

Lemma 4.17. $\langle I(H_2), S_{\psi_\delta} \rangle = O(\delta)$.

Proof. Divide $\langle I(H_2), S_\psi \rangle$ into two integrations; the integration over

$$R_{\geq 1} := \{(x; y) \in C_{H_2}(S_\psi) \,|\, y \notin \mathbb{R}^{4k} \times (-1,1)^{2k}\}$$

($x = (x_1, x_2, x_3)$) and that over $R_{<1} := C_{H_2}(S_\psi) \setminus R_{\geq 1}$.

The integration over $R_{\geq 1}$ is well defined and continuous at $\delta = 0$, since $y \in \mathbb{R}^{6k}$ is far from the image of S_{ψ_δ} (see (4.8)). But when $\delta = 0$, this integral is zero, since all the three points $S_\psi(x_p)$ ($p = 1,2,3$) are in $\mathbb{R}^{4k} \times \{0\}^{2k}$ and hence the image of the direction map $\varphi_{H_2} := \varphi^\theta_{14} \times \varphi^\theta_{24} \times \varphi^\theta_{34}$ is of positive codimension $\geq 2k-1$ in $(S^{6k-1})^3$ (see Lemma 4.4). Hence the integration over $R_{\geq 1}$ is $O(\delta)$.

Next consider the integration over $R_{<1}$. We have only to consider $(x; y)$ with the first $6k-1$ factors of $S_\psi(x_1), S_\psi(x_2), S_\psi(x_3)$ and y close to each other; otherwise the image of φ_{H_2} cannot be in $(\mathrm{supp}(vol))^3$ (see condition (2) just before Figure 4.2). In particular all $(s^{(p)}, t^{(p)}) \in \mathbb{R}^{4k-2}$ ($p = 1,2,3$) must be close to each other.

Consider the case that $x = (x_1, x_2, x_3) \in C_3(\mathbb{R}^{4k-1} \setminus L)$. Similarly as in Lemma 4.12, if we choose $vol_{S^{6k-1}}$ with sufficiently small support, then we may assume that at most one of $x \cap (X \setminus L)$ and $x \cap (Y \setminus L)$ is non-empty. We can prove the vanishing of the integration of ω_{H_2} over $\{(x;y) \,|\, x \cap X \neq \emptyset\}$ in a similar way as in Lemma 4.12 by considering an involution $F_1 : (x_1, x_2, x_3; y) \mapsto (i_1 x_1, i_1 x_2, i_1 x_3; i_1 y)$ (i_1 was defined in Lemma 4.12), which preserves the orientation but satisfies $F_1^* \omega_{H_2} = -\omega_{H_2}$ because $\varphi^\theta_{p4} \circ F_1 = i_1 \circ \varphi^\theta_{p4}$ ($p = 1,2,3$) on X by (4.9) and $i_1^* vol_{S^{6k-1}} = -vol_{S^{6k-1}}$. The vanishing of the integrations over $\{(x;y) \,|\, x \cap Y \neq \emptyset\}$ and $\{x \cap X = x \cap Y = \emptyset\}$ can also be proved by similar involutions F_i, $i = 1, 2$.

So we may assume that one of x_p is in L (and other two points are near L). Since the first $6k-1$ factors of $S_\psi(x_p)$ ($p = 1,2,3$) are close to each other, an analogous argument to the proof of Lemma 4.4 shows that only the integration over the subspace of $(x;y)$ with x in $C_3(N_1 \sqcup N_3)$ or $C_3(N_2 \sqcup N_4)$ or $C_3(N'_1 \sqcup N'_2)$, where $N'_i := \mathbb{R}^{4k-2} \times (\zeta_i - \varepsilon, \zeta_i + \varepsilon)$ ($i = 1,2$) correspond to the crossing c of f (see §4.2.2 and Figure 4.3); otherwise two or more $S_\psi(x_p)$'s are in $\mathbb{R}^{4k} \times \{0\}^{2k}$ and hence the image of the map φ_{H_2} is of positive codimension $\geq 2k-1$ in $(S^{6k-1})^3$. But similarly as in Lemma 4.16, on these spaces we can define translations τ, τ' under which φ_{H_2} is invariant, and hence the integrand ω_{H_2} must vanish by dimensional reason. $\square$

Contribution of the correction term c. Lastly we compute the third term of (4.7). This is an integration of $D(S_\psi)^*\mu$ over $\mathbb{R}^{4k-1}$ (cf. Lemma 4.5). See Lemma 5.22 and Definition 5.23 for the definition of c.

Lemma 4.18. *$\langle c,\, S_\psi\rangle = 0$.*

Proof. First we show that integrations of $D(S_\psi)^*\mu$ over X, Y and Z (see §4.2.2) vanish. For X, this is because

- $i_1^* D(S_\psi)^*\mu = D(S_\psi)^*\mu$ on X, since $D(S_\psi)\circ i_1 = i_1 \circ D(S_\psi)$ on X by (4.9) (where $i_1 : \mathcal{I}_{4k-1}(\mathbb{R}^{6k}) \to \mathcal{I}_{4k-1}(\mathbb{R}^{6k})$ is given by $f \mapsto i_1 \circ f$; see Remark 5.24) and we can choose μ so that $i_1^*\mu = \mu$ (see Remark 5.24), and
- i_1 is an orientation reversing diffeomorphism of X.

Similar arguments hold for Y and Z.

So we may restrict the integration to $[-1/2, 1/2]^{4k-2}\times\mathbb{R}^1$. If $(s,t,u)\notin N_3 \sqcup N_4$, then $\psi(s,t)(u)$ does not depend on $(s,t)\in\mathbb{R}^{4k-2}$, so $D(S_\psi)$ is invariant under the translations τ and τ' defined in the proof of Lemma 4.16. If $(s,t,u)\in N_3$ (resp. N_4), then $\psi(s,t)(u)$ does not depend on $t \in \mathbb{R}^{2k-1}$ (resp. s), so $D(S_\psi)$ is invariant under the translations τ (resp. τ'). Thus the image of $[-1/2,1/2]^{4k-2}\times\mathbb{R}^1$ via $D(S_\psi)$ must be of dimension $< 4k-1$ and hence a $(4k-1)$-form $D(S_\psi)^*\mu$ must vanish on $[-1/2,1/2]^{4k-2}\times\mathbb{R}^1$. $\square$

Thus we have completed the proof of Theorem 4.7; only Type II configurations for H_1 contribute to $\langle\mathcal{H},\, S_\psi\rangle$ by ±1, and hence $\langle\mathcal{H},\, S_\psi\rangle = \pm 1$.

4.3. **Non-triviality of $\mathcal{H}$ in general dimensions.**

Here we complete the proof of Theorem 1.3. Suppose that $n > j \geq 2$, $n-j\geq 3$ is odd and $m := 2n-3j-3 > 0$. Put $n - j = 2k+1$ ($k \geq 1$) and consider $S_\psi \in \mathcal{K}_{6k,4k-1}$ as above. Notice that $n = 6k-m$ and $j = 4k-m-1$, and in particular $4k-m-1>0$.

Since S_ψ is of the form (4.6), we can find $l_m \in \Omega^m\mathcal{K}_{6k-m,4k-m-1} = \Omega^m\mathcal{K}_{n,j}$ such that $S_\psi = \mathrm{gr}^m(l_m)$. Explicitly we can define $l_m : \mathbb{R}^m \to \mathcal{K}_{n,j}$ by

$$\begin{aligned} &l_m(t_1,\dots,t_m)(x_1,\dots,x_j) := \\ &\qquad ((x_1,\dots,x_{j-1}), \psi(t_1,\dots,t_m,x_1,\dots,x_{j-1})(x_j)) \in \mathbb{R}^{j-1}\times\mathbb{R}^{2k+2} = \mathbb{R}^n \end{aligned} \tag{4.12}$$

and regard it as in $\Omega^m\mathcal{K}_{n,j}$. We think of $[l_m]$ as the generator of $H_m(\mathcal{K}_{n,j})$ via the Hurewicz isomorphism ($\mathcal{K}_{n,j}$ is $(2n-3j-4)$-connected; see [4]).

Consider $\mathcal{H} = [I(H)+c] \in H^m_{DR}(\mathcal{K}_{n,j})$. The following completes the proof of Theorem 1.3.

Theorem 4.19. *The Kronecker pairing $\langle\mathcal{H},\, l_m\rangle$ is equal to ±1.*

Proof. Define the spaces $\hat{C}_{H_i}$, $i = 1,2$, by $\hat{C}_{H_1} := \mathbb{R}^m \times C_4^o(\mathbb{R}^j)$ and

$$\hat{C}_{H_2} := \{(t,(x_1,x_2,x_3),y) \in \mathbb{R}^m\times C_3^o(\mathbb{R}^j)\times\mathbb{R}^n \mid l_m(t)(x_p) \neq y,\ p=1,2,3\}.$$

These spaces are also defined as the following pullback square;

$$\begin{tikzcd} \hat{C}_{H_i} \arrow[r, "\hat{l}_m"] \arrow[d] & C_{H_i} \arrow[d, "\pi_{H_i}"] \\ \mathbb{R}^m \arrow[r, "l_m"] & \mathcal{K}_{n,j} \end{tikzcd}$$

Then $\langle \mathcal{H},\, l_m\rangle$ is equal to

$$\frac{1}{4}\int_{\hat{C}_{H_1}} \hat{l}_m^* \omega_{H_1} + \frac{1}{12}\int_{\hat{C}_{H_2}} \hat{l}_m^* \omega_{H_2} + \int_{\mathbb{R}^m\times\mathbb{R}^j} D(l_m)^*\mu, \tag{4.13}$$

where $D(l_m) : \mathbb{R}^m \times \mathbb{R}^j \to \mathcal{I}_j(\mathbb{R}^n)$ is defined by $(t, x) \mapsto d(l_m(t))_x$.

Recall $C_{H_1}(S_\psi) = C_4^o(\mathbb{R}^{4k-1})$ and $C_{H_2}(S_\psi) \subset C_3^o(\mathbb{R}^{4k-1}) \times \mathbb{R}^{6k}$. We regard $\hat{C}_{H_i} \subset C_{H_i}(S_\psi)$ $(i = 1, 2)$;

$$\hat{C}_{H_1} \cong \{(x_1, \dots, x_4) \in C_{H_1}(S_\psi) \,|\, pr_m(x_1) = \cdots = pr_m(x_4)\},$$

$$\hat{C}_{H_2} \cong \{(x_1, x_2, x_3; y) \in C_{H_2}(S_\psi) \,|\, pr_m(x_1) = pr_m(x_2) = pr_m(x_3) = pr_m(y)\}$$

($pr_m : \mathbb{R}^N \to \mathbb{R}^m$ ($N = 4k-1$ or $6k$) is the first m projection), via diffeomorphisms given by respectively

$$(t, x) \longmapsto ((t, x_1), \dots, (t, x_4)), \quad (t, x, y) \longmapsto ((t, x_1), (t, x_2), (t, x_3), (t, y)).$$

The direction maps $C_{H_i} \to S^{N-1}$, $N = n$ or j, composed by $\hat{l}_m$ are regarded as

$$\begin{aligned} &\varphi^\theta_{12} : \hat{C}_{H_1} \longrightarrow S^{n-1} = S^{6k-1} \cap \{x \in \mathbb{R}^{6k}\,;\, pr_m(x) = 0\},\\ &\varphi^\theta_{12}((t, x_1), (t, x_2)) = \frac{(x_2 - x_1, S_\psi(t, x_2) - S_\psi(t, x_1))}{|(x_2 - x_1, S_\psi(t, x_2) - S_\psi(t, x_1))|}, \end{aligned} \tag{4.14}$$

and so on. Then the integrations relating to $L \subset \mathbb{R}^m \times \mathbb{R}^j = \mathbb{R}^{4k-1}$ (a neighborhood of $[-1/2, 1/2]^m \times [-1/2, 1/2]^{j-1} \times \mathbb{R}^1$) can be computed in similar ways as in the previous subsection;

The first term. Type I contribution vanishes by the translations τ or τ' as in Lemma 4.16. Type II configurations contribute by ±2; each component

$$(N^{l,m}_{i,i+2,i+1,i+3}) \cap \{pr_m(x_1) = \cdots = pr_m(x_4)\} \quad \text{or}$$

$$(N^{l,m}_{i,i+2,i+3,i+1}) \cap \{pr_m(x_1) = \cdots = pr_m(x_4)\}$$

for some $i = 1, \dots, 4$ and $l, m = \pm$ is mapped via the direction map to some

$$U^{6k-1}_l \times U^{6k-1}_{l'} \times U^{4k-2}_{l''} \cap \{\text{first } m \text{ projections} = 0\},$$

which is $U^{n-1}_l \times U^{n-1}_{l'} \times U^{j-1}_{l''}$ (see §4.2.3). The sign arguments are slightly different, but the result is same as Lemma 4.15; in the first diagram in the proof of Lemma 4.15, the map G restricted to $\hat{C}_{H_1}$ preserve the orientation but the left vertical map does not. Both the vertical maps in the second diagram restricted to $\hat{C}_{H_1}$ have the same orientation sign $(-1)^a$, where $a = \min\{m, 2k-1\}$.

The second term. Similarly as in Lemma 4.17, the integration over $R_{\geq 1} := \{(t, x, y) \in \hat{C}_{H_2} \,|\, y \notin \mathbb{R}^{j+1} \times (-1, 1)^{2k}\}$ (notice $j + 1 + 2k = n$) is $O(\delta)$. In $R_{<1} := \hat{C}_{H_2} \setminus R_{\geq 1}$, only (t, x, y) with $pr_{j-1}(x_p)$ $(p = 1, 2, 3)$ close to each other and $x \in C_3((N_i \sqcup N_{i+2}) \cap L)$ $(i = 1, 2)$ or $C_3((N'_1 \sqcup N'_2) \cap L)$ may contribute to the integral (where $N_i := \mathbb{R}^m \times \mathbb{R}^{j-1} \times (\xi_i - \varepsilon, \xi_i + \varepsilon)$, $N'_i := \mathbb{R}^m \times \mathbb{R}^{j-1} \times (\zeta_i - \varepsilon, \zeta_i + \varepsilon)$); otherwise the image of φ_{H_2} is of positive codimension. But on these spaces the direction map is invariant under the similar translations τ or τ' to those in Lemma 4.16 and the integrand vanishes by dimensional reason.

The third term. Similarly as in Lemma 4.18, since the derivation map $D(l_m)$ is invariant on L under the translations given by using τ or τ', the integration over L vanishes by dimensional reason.

The signs appearing in the proof of vanishing of integrations over $\mathbb{R}^{4k-1} \setminus L$ are slightly different from those in the previous subsection. Recall the involutions F_1 and F_2 on $C_{H_i}(S_\psi)$. They clearly preserve $\hat{C}_{H_i}$, and are given by

$$\begin{aligned} F_l((t,x_1),\dots,(t,x_4)) &= (i_l(t,x_1),\dots,i_l(t,x_4)) && \text{on } \hat{C}_{H_1}, \\ F_l((t,x_1),(t,x_2),(t,x_3);(t,y)) &= (i_l(t,x_1),i_l(t,x_2),i_l(t,x_3);i_l(t,y)) && \text{on } \hat{C}_{H_2} \end{aligned}$$

($l = 1, 2$), here we regard $\hat{C}_{H_i}$ as a subspace of $C_{H_i}(S_\psi)$ as above. The maps $i_1, i_2 : \mathbb{R}^N \to \mathbb{R}^N$ ($N = 4k-1$ or $6k$) are as given in Lemma 4.12. For $N = 4k-1$, they are written explicitly as

$$i_1(t,x) = \begin{cases} (-t_1,\dots,-t_m; -x_1,\dots,-x_{2k-1-m}, x_{2k-m},\dots,x_j) & m \le 2k-1, \\ (-t_1,\dots,-t_{2k-1}, t_{2k},\dots,t_m; x_1,\dots,x_j) & m > 2k-1, \end{cases}$$

$$i_2(t,x) = \begin{cases} (t_1,\dots,t_m; x_1,\dots,x_{2k-1-m}, -x_{2k-m},\dots,-x_{j-1},x_j) & m \le 2k-1, \\ (t_1,\dots,t_{2k-1}, -t_{2k},\dots,-t_m; -x_1, \cdots - x_{j-1}, x_j) & m > 2k-1. \end{cases}$$

First consider the first and the second terms of (4.13). Let $X, Y, Z \subset \mathbb{R}^m \times \mathbb{R}^j = \mathbb{R}^{4k-1}$ be subsets defined similarly as in §4.2.2, and set

$$\begin{aligned} \hat{C}_{H_1}(X) &:= \{(t,(x_1,\dots,x_4)) \in \hat{C}_{H_1} \mid (t,x_p) \in X, \ \forall p\}, \\ \hat{C}_{H_2}(X) &:= \{(t,(x_1,x_2,x_3),y) \in \hat{C}_{H_2} \mid (t,x_p) \in X, \ \forall p\} \end{aligned}$$

and so on. Then the actions of F_1 and F_2 on the forms ω_{H_i} and orientations of the spaces are described as in Table 4.1, which is a consequence of the equations

$$\varphi \circ F_1 = \begin{cases} i_{1,2k-m-1} \circ \varphi & m \le 2k-1 \\ \varphi & m > 2k-1 \end{cases} \quad \text{on } \hat{C}_{H_i}(X),$$

$$\varphi \circ F_2 = \begin{cases} i_{2k-m,j-1} \circ \varphi & m \le 2k-1 \\ i_{1,j-1} \circ \varphi & m > 2k-1 \end{cases} \quad \text{on } \hat{C}_{H_i}(Y),$$

where φ is one of the direction maps, and $i_{p,q} : \mathbb{R}^n \to \mathbb{R}^n$ ($p < q$) is given by

$$i_{p,q}(a_1,\dots,a_n) := (a_1,\dots,a_{p-1},-a_p,\dots,-a_q,a_{q+1},\dots,a_n)$$

(in particular i_1 and i_2 we have used can be written as $i_1 = i_{1,2k-1}$, $i_2 = i_{2k,4k-2}$). Thus the integrations of ω_{H_i} over $\hat{C}_{H_i}(X)$ and $\hat{C}_{H_i}(Y)$ vanish (when $m > 2k-1$, we use $j = 4k-1-m$ and hence $m - 2k + j$ is odd).

	$m \le 2k-1$	$m > 2k-1$
orientation sign of F_1	$(-1)^m$	-1
$F_1^*\omega_{H_i}$ on $\hat{C}_{H_i}(X)$	$(-1)^{2k-1-m}\omega_{H_i}$	$+\omega_{H_i}$
orientation sign of F_2	$+1$	$(-1)^{m-2k+1}$
$F_2^*\omega_{H_i}$ on $\hat{C}_{H_i}(Y)$	$-\omega_{H_i}$	$(-1)^{j-1}\omega_{H_i}$

Table 4.1. Signs of F_l

For the third term of (4.13), we have to study the signs arising from the involutions i_1 and i_2. The orientation signs of $i_1, i_2 : \mathbb{R}^m \times \mathbb{R}^j \to \mathbb{R}^m \times \mathbb{R}^j$ are always

-1. But i_1, i_2 always preserve the integrand, because

$$D(l_m) \circ i_1 = \begin{cases} i_{1,2k-m-1} \circ D(l_m) & \text{if } m \leq 2k-1, \\ D(l_m) & \text{if } m > 2k-1, \end{cases} \quad \text{on } X \cup Z,$$

$$D(l_m) \circ i_2 = \begin{cases} i_{2k-m,j-1} \circ D(l_m) & \text{if } m \leq 2k-1, \\ i_{1,j-1} \circ D(l_m) & \text{if } m > 2k-1, \end{cases} \quad \text{on } Y \cup Z,$$

and we can choose μ so that i_1', i_2' and i_2'' preserve μ (see Remark 5.24). Hence the integrations of μ over X, Y and Z vanish. □

5. Vanishing results

In §§5.1, ..., 5.8 we prove Theorem 3.4 assuming $n-j \geq 2$ is even, by studying the boundary strata of compactified configuration spaces. Some results here hold even if $n-j \geq 3$ is odd and can be used to prove Theorem 4.1 (see §5.9). In §5.10 we complete the proof of Proposition 3.5.

5.1. **Boundary strata.** Let Γ be a graph with s i-vertices and t e-vertices. We denote by $C_{s,t}$ the fiber of π_Γ. The compactified configuration space $C_{s,t}$ is a manifold with corners. The boundary $\partial C_{s,t}$ consists of configurations where some points in the configuration are allowed to 'collide together.' Moreover $\partial C_{s,t}$ is stratified via 'complexities' of collisions.

But here we do not need the complete description of all strata. For our purpose only the most 'generic' part, the codimension one strata, are needed. Such strata correspond to 'coinstantaneous collisions' of points, and are parametrized by subsets of the set $V(\Gamma)$ of vertices of Γ, as we will explain in §5.2.

5.2. **Codimension one strata.** To any subset $A \subset V(\Gamma)$ with $\sharp A \geq 2$, a codimension one stratum $C_A \subset \partial C_{s,t}$ is assigned. Namely C_A consists of configurations where the points labeled by A 'simultaneously collide together.' More precisely, any point in C_A can be written as a limit point

$$\lim_{\tau \to 0} (x_1(\tau), \dots, x_s(\tau), y_{s+1}(\tau), \dots, y_{s+t}(\tau)) \tag{5.1}$$

such that $(x_1(\tau), \dots, y_{s+t}(\tau)) \in C^o_{s,t}$ for $\tau > 0$, and $x_p(\tau)$, $y_q(\tau)$ can be written as

$$x_p(\tau) = \begin{cases} x_p \quad (\text{constant}) & \text{if } p \notin A, \\ z + \tau v_p & \text{if } p \in A, \end{cases} \qquad y_q(\tau) = \begin{cases} y_q \quad (\text{constant}) & \text{if } q \notin A, \\ z + \tau w_q & \text{if } q \in A, \end{cases}$$

for some $v_p \in \mathbb{R}^j \setminus \{0\}$, $w_q \in \mathbb{R}^n \setminus \{0\}$ and $z \in \mathbb{R}^n$.

There are other types of codimension one strata, denoted by C^∞_A, parametrized by all the non-empty subsets $A \subset V(\Gamma)$. The stratum $C^\infty_A \subset \partial C_{s,t}$ consists of configurations where the points labeled by A 'escape to infinity.' More precisely C^∞_A is the set of limit points

$$\lim_{\tau \to \infty} (x_1(\tau), \dots, x_s(\tau), y_{s+1}(\tau), \dots, y_{s+t}(\tau)) \tag{5.2}$$

where $(x_1(\tau), \dots, y_{s+t}(\tau)) \in C^o_{s,t}$ $(0 < \tau < \infty)$ is of the form

$$x_p(\tau) = \begin{cases} x_p \quad (\text{constant}) & \text{if } p \notin A, \\ x_p + \tau v_p & \text{if } p \in A, \end{cases} \qquad y_q(\tau) = \begin{cases} y_q \quad (\text{constant}) & \text{if } q \notin A, \\ y_q + \tau w_q & \text{if } q \in A, \end{cases}$$

for some $v_p \in \mathbb{R}^j \setminus \{0\}$, $w_q \in \mathbb{R}^n \setminus \{0\}$, $x_p \in \mathbb{R}^j$ and $y_q \in \mathbb{R}^n$.

All the codimension one strata is of the form C_A or C_A^∞, hence we have

$$\partial C_{s,t} = \left(\bigcup_{A\subset V(\Gamma),\ \sharp A\geq 2} \overline{C_A} \right) \cup \left(\bigcup_{A'\subset V(\Gamma),\ \sharp A'\geq 1} \overline{C^\infty_{A'}} \right).$$

We will call C_A ($\sharp A \geq 2$) a *stratum of non-infinity type*, and C_A^∞ ($A \neq \emptyset$) a *stratum at infinity*.

Define the subsets Σ_A and Σ_A^∞ of C_Γ by

$$\Sigma_A := \bigcup_{f\in\mathcal{K}_{n,j}} C_A(f), \quad \Sigma_A^\infty := \bigcup_{f\in\mathcal{K}_{n,j}} C_A^\infty(f)$$

where $C_A(f)$ and $C_A^\infty(f)$ are codimension one strata of $\pi_\Gamma^{-1}(f)$ described as above. Then Σ_A and Σ_A^∞ fiber over $\mathcal{K}_{n,j}$.

Notice that it is enough to describe $\mathrm{Int}\,\Sigma_A^{(\infty)}$ for the proof of Theorem 3.4. In §5.3 and §5.4 we will describe these strata explicitly, following [3, 5, 14, 20].

5.3. **Explicit description of non-infinity type strata.** Let $A \subset V(\Gamma)$ be a subset with $\sharp A \geq 2$ (recall that $V(\Gamma)$ denotes the set of vertices of a graph Γ). Here we study the strata Σ_A of non-infinity type. Denote by $E(\Gamma)$ the set of edges of Γ.

Definition 5.1. The *subgraph* Γ_A of Γ associated with A is a (possibly non-admissible) graph with $V(\Gamma_A) = A$ and $E(\Gamma_A) = \{pq \in E(\Gamma) \,|\, p, q \in A,\ p \neq q\}$ (hence small loops are ignored). If $A = \{p_1 < \cdots < p_k\}$, then the vertex of Γ_A which was labeled by p_a in Γ is re-labeled by a. The labels of edges are suitably decreased.

The *quotient graph* Γ/Γ_A is a graph obtained by 'collapsing Γ_A to a point v_A.' More precisely,

$$V(\Gamma/\Gamma_A) := (V(\Gamma) \setminus A) \sqcup \{v_A\},$$
$$E(\Gamma/\Gamma_A) := \{pq \in E(\Gamma) \,|\, p, q \notin A\} \sqcup \{pv_A \,|\, p \notin A,\ pq \in E(\Gamma) \text{ for } \exists q \in A\}.$$

The vertex v_A is internal if there is an i-vertex in A, and is external otherwise. We label v_A by $\min\{p\,;\, p \in A\}$, and the labels of other vertices and edges are suitably decreased (see Figure 5.1 for an example).



FIGURE 5.1. Examples of Γ_A and Γ/Γ_A for $A = \{1, 4\}$ (n odd)

There is a projection

$$p_A : \mathrm{Int}\,\Sigma_A \longrightarrow C^o_{\Gamma/\Gamma_A}$$

which maps the limit point (5.1) to $(\ldots, x_p, \ldots, z, \ldots, y_q, \ldots)$, $p, q \notin A$. The point z corresponds to the vertex v_A, hence $z \in \mathbb{R}^n$ if A contains no i-vertex, and $z \in \mathbb{R}^j$ if there is an i-vertex in A.

The fiber of p_A is thought of as the space of 'infinitesimal configurations' at the colliding point. We will define a fibration $\rho_A : \hat{B}_A \to B_A$ which describes such infinitesimal configurations.

Definition 5.2. Define $\mathcal{I}_j(\mathbb{R}^n)$ as the set of all j-frames in $\mathbb{R}^n$. In other words

$$\mathcal{I}_j(\mathbb{R}^n) := \{\text{linear injective maps } \mathbb{R}^j \hookrightarrow \mathbb{R}^n\}.$$

We give $\mathcal{I}_j(\mathbb{R}^n)$ a natural structure of an open submanifold of $(\mathbb{R}^n \setminus \{0\})^j$.

Let a and b be the numbers of i- and e-vertices in A. Define a manifold B_A by

$$B_A := \begin{cases} \{*\} & a = 0, \\ \mathcal{I}_j(\mathbb{R}^n) & a > 0. \end{cases}$$

When $a = 0$, we define $\hat{B}_A$ to be $C^o_{\sharp A}(\mathbb{R}^n)$ modulo scaling and translation;

$$\hat{B}_A := C^o_b(\mathbb{R}^n)/\mathbb{R}^n \rtimes \mathbb{R}_{>0},$$

where the action of $\mathbb{R}^n \rtimes \mathbb{R}_{>0}$ is defined by

$$(y_p)_{p\in A} \longmapsto (\alpha(y_p - \beta))_{p\in A}, \quad \forall \alpha > 0,\ \forall \beta \in \mathbb{R}^n.$$

The map $\rho_A : \hat{B}_A \to B_A = \{*\}$ is defined as the canonical one.

When $a > 0$, $\hat{B}_A$ is $\mathcal{I}_j(\mathbb{R}^n) \times C^o_{a,b}$ modulo scaling and translation in the directions of j-planes;

$$\hat{B}_A := (\mathcal{I}_j(\mathbb{R}^n) \times C^o_{a,b})/\mathbb{R}^j \rtimes \mathbb{R}_{>0},$$

where the action of $\mathbb{R}^j \rtimes \mathbb{R}_{>0}$ is defined by

$$(\iota; (x_p, y_q)_{p,q\in A}) \longmapsto (\iota; (\alpha(x_p - \beta), \alpha(y_q - \iota(\beta)))_{p,q\in A}), \quad \forall \alpha > 0,\ \forall \beta \in \mathbb{R}^j.$$

The map $\rho_A : \hat{B}_A \to B_A = \mathcal{I}_j(\mathbb{R}^n)$ is defined as the natural projection.

Finally define $D_A : C^o_{\Gamma/\Gamma_A} \to B_A$ to be the canonical map if $a = 0$, and

$$D_A(f; \dots, x_p, \dots, z, \dots, y_q, \dots)_{p,q\notin A}) := (df_z : T_z\mathbb{R}^j \longrightarrow T_{f(z)}\mathbb{R}^n) \in \mathcal{I}_j(\mathbb{R}^n)$$

if $a > 0$ (each tangent spaces are naturally identified with $\mathbb{R}^j$ and $\mathbb{R}^n$).

Proposition 5.3 ([3, 5, 14, 20])**.** *The fibration* $p_A : \operatorname{Int}\Sigma_A \to C^o_{\Gamma/\Gamma_A}$ *is the pull-back of* ρ_A *via* D_A*;*

$$\begin{tikzcd}
\operatorname{Int}\Sigma_A \arrow[r, "\hat{D}_A"] \arrow[d, "p_A"'] & \hat{B}_A \arrow[d, "\rho_A"] \\
C^o_{\Gamma/\Gamma_A} \arrow[r, "D_A"] & B_A
\end{tikzcd}$$

In particular $\operatorname{Int}\Sigma_A \approx C^o_{\Gamma/\Gamma_A} \times \hat{B}_A$ *if* A *has no i-vertex.*

Notice that the differential form $\omega_{\Gamma/\Gamma_A} \in \Omega^*_{DR}(C^o_{\Gamma/\Gamma_A})$ can be defined similarly to §3.2, by using the direction maps φ corresponding to the edges of Γ/Γ_A. Similarly, the maps φ_e for any edges of Γ_A are well defined on $\hat{B}_A$; if $e = \overrightarrow{pq}$ is an η-edge,

$$\hat{\varphi}^\eta_e(\iota; (x_r, y_s)_{r,s\in A}) := \frac{x_q - x_p}{|x_q - x_p|} \in S^{j-1},$$

and if e is a θ-edge,

$$\hat{\varphi}^\theta_e(\iota; (x_r, y_s)_{r,s\in A}) := \frac{z_q - z_p}{|z_q - z_p|} \in S^{n-1},$$

where $z_p = x_p$ or $\iota(y_p)$ according to whether p is internal or external. Hence

$$\hat{\omega}_{\Gamma_A} := \bigwedge_{e\in E(\Gamma_A)} \varphi^*_e vol \in \Omega^*_{DR}(\hat{B}_A)$$

can be defined. Then we have

$$\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = \pm(p_A^*\omega_{\Gamma/\Gamma_A}) \wedge (\hat{D}_A^*\hat{\omega}_{\Gamma_A})$$

and hence

$$(p_A)_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = \pm\omega_{\Gamma/\Gamma_A} \wedge D_A^*(\rho_A)_*\hat{\omega}_{\Gamma_A} \tag{5.3}$$

by the compatibility of fiber-integrations with pullbacks (for signs see §§5.5, 5.6).

Denote $\pi_\Gamma^{\partial_A} := \pi_\Gamma|_{\mathrm{Int}\,\Sigma_A} : \mathrm{Int}\,\Sigma_A \to \mathcal{K}_{n,j}$. Notice that $\pi_\Gamma^{\partial_A} = \pi_{\Gamma/\Gamma_A} \circ p_A$. We will often use the following criterion to show the vanishing of an integration along $\mathrm{Int}\,\Sigma_A$.

Lemma 5.4 ([5]). *Let a and b be the numbers of i- and e-vertices in A respectively. Then the fiber integration $(\pi_\Gamma^{\partial_A})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A}$ vanishes unless*

$$\begin{aligned} &\deg\hat{\omega}_{\Gamma_A} = nb - (n+1) && \text{if } a = 0,\\ &0 \le \deg\hat{\omega}_{\Gamma_A} - (ja + nb - (j+1)) \le nj && \text{if } a > 0. \end{aligned}$$

Proof. The form $(\rho_A)_*\hat{\omega}_{\Gamma_A}$ vanishes unless $0 \le \deg(\rho_A)_*\hat{\omega}_{\Gamma_A} \le \dim B_A$. We have

$$\deg(\rho_A)_*\hat{\omega}_{\Gamma_A} = \deg\hat{\omega}_{\Gamma_A} - \dim(\text{fiber of } \rho_A : \hat{B}_A \to B_A)$$

and, by definition of the fibration $\rho_A : \hat{B}_A \to B_A$,

$$(\dim B_A,\ \dim(\text{fiber of } \rho_A)) = \begin{cases} (0,\ nb - (n+1)) & \text{if } a = 0,\\ (nj,\ ja + nb - (j+1)) & \text{if } a > 0. \end{cases}$$

Hence $(\rho_A)_*\hat{\omega}_{\Gamma_A}$ vanishes unless the conditions of the Lemma are satisfied. Then the formulas $(\pi_\Gamma^{\partial_A})_* = \pm(\pi_{\Gamma/\Gamma_A})_* \circ (p_A)_*$ and (5.3) complete the proof. □

5.4. **Explicit description of strata at infinity.** Let Γ be a graph and A a non-empty subset of $V(\Gamma)$. Below we describe $\mathrm{Int}\,\Sigma_A^\infty$ following [3, 5, 14, 20].

Definition 5.5. Define the *complementary graph* Γ_A^c by letting

$$V(\Gamma_A^c) := V(\Gamma) \setminus A, \quad E(\Gamma_A^c) := \{\overrightarrow{pq} \in E(\Gamma) \,|\, p, q \notin A\}$$

($\Gamma_A^c := \emptyset$ if $A = V(\Gamma)$). A *graph with infinity* is a graph with a specified vertex v^∞ (called *vertex at infinity*), which is not regarded as being internal nor external.

Define Γ_A^∞, a graph with infinity, by 'shrinking Γ_A^c to a point.' Namely Γ_A^∞ is defined similarly as a quotient graph Γ/Γ_A^c (see Definition 5.1), but its vertex v_A is replaced by v^∞, a vertex at infinity (see Figure 5.2 for an example). By definition $\Gamma_A^\infty := \Gamma$ if $A = V(\Gamma)$.



FIGURE 5.2. Examples of Γ_A^c and Γ_A^∞ for $A = \{3, 4\}$ (n even)

$\mathrm{Int}\,\Sigma_A^\infty$ fibers over $C^o_{\Gamma_A^c}$;

$$p_A^\infty : \mathrm{Int}\,\Sigma_A^\infty \longrightarrow C^o_{\Gamma_A^c}$$

which maps the limit point (5.2) in §5.2 to $(x_p, y_q)_{p,q\notin A}$. In other words p_A^∞ forgets the points escaping to infinity.

As in §5.3, we will define a space $\hat{B}_A^\infty$ which describes infinitesimal configurations around infinity. The space $\hat{B}_A^\infty$ is a subquotient of $C_{a,b}^o$ (a, b are the numbers of i- and e-vertices in A) modulo scaling;

$$\hat{B}_A^\infty := \{(x_p, y_q)_{p,q\in A} \in C_{a+b}^o(\mathbb{R}^n \setminus 0) \,|\, x_p \in \mathbb{R}^j \times \{0\}^{n-j}\}/\mathbb{R}_{>0}.$$

The origin $0 \in \mathbb{R}^n$ corresponds to v^∞, which we use to fix the coordinates. So in this case translation is not needed (compare it with the definition of $\hat{B}_A$).

Proposition 5.6 ([3, 5, 14, 20])**.** $\mathrm{Int}\,\Sigma_A^\infty$ *is homeomorphic to* $C_{\Gamma_A^c}^o \times \hat{B}_A^\infty$.

The form $\omega_{\Gamma_A^\infty} \in \Omega_{DR}^*(\hat{B}_A^\infty)$ can be defined as in §3.2 since the direction maps $\hat{\varphi}$ are invariant under scaling. The vertex v^∞ corresponds to $0 \in \mathbb{R}^n$. More precisely, for each $e \in E(\Gamma_A^\infty)$, define the maps $\varphi_e : \hat{B}_A^\infty \to S^{N-1}$ ($N = j$ or n) by

$$\varphi_e(x_p, y_q)_{p,q\in A} := \begin{cases} (z_q - z_p)/|z_q - z_p| & e = \overrightarrow{pq},\ p, q \in A, \\ -z_p/|z_p| & e = \overrightarrow{pv^\infty},\ p \in A, \end{cases}$$

where z_p denotes x_p or y_p according to whether p is an i-vertex or not. Then

$$\omega_{\Gamma_A^\infty} := \bigwedge_{e\in E(\Gamma_A^\infty)} \omega_e.$$

Under the identification in Proposition 5.6,

$$\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A^\infty} = \pm pr_1^*\omega_{\Gamma_A^c} \wedge pr_2^*\omega_{\Gamma_A^\infty}.$$

Define $\pi_\Gamma^{\partial_A^\infty} : \mathrm{Int}\,\Sigma_A^\infty \to \mathcal{K}_{n,j}$ as the restriction of π_Γ onto the stratum $\mathrm{Int}\,\Sigma_A^\infty$ of fibers. Then we have the following (which should be compared with Lemma 5.4).

Lemma 5.7 ([5])**.** *Let* a *and* b *be the numbers of i- and e-vertices of* Γ_A^∞ *respectively (other than* v^∞*). Then the integration* $(\pi_\Gamma^{\partial_A^\infty})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A^\infty}$ *vanishes unless* $\deg\omega_{\Gamma_A^\infty} = \dim\hat{B}_A^\infty$*, or equivalently, unless*

$$\deg\omega_{\Gamma_A^\infty} = ja + nb - 1.$$

5.5. **Orientations of boundary strata.** Let Γ be a graph and A a non-empty subset of $V(\Gamma)$.

Definition 5.8. The boundary face Σ_A (or the subgraph Γ_A) is said to be *principal* if A consists of exactly two vertices. Similarly the boundary face Σ_A^∞ is said to be *principal* if A consists of exactly one vertex.

All the other boundary strata are said to be *hidden*.

Here we study the induced orientations of the non-infinity type principal strata C_A from that of $\mathrm{fib}(\pi_\Gamma) \approx C_{s,t}$ (see §5.2). We are not interested in the orientations of strata at infinity and hidden strata, since the integrations along these strata will be proved to vanish (see below).

Let s and t be the numbers of i- and e-vertices of Γ respectively. The fiber $C_{s,t}$ is equipped with the natural orientation as the subspace of $(\mathbb{R}^j)^s \times (\mathbb{R}^n)^t$.

Let $A \subset V(\Gamma)$ be a subset with $\sharp A = 2$ (thus Σ_A is principal), and a and b (with $a + b = 2$) the numbers of i- and e-vertices in A.

Case 1: $a = 2$, $b = 0$.

Let $A = \{x_p, x_q\}$ $(p < q \leq s)$. In this case $\hat{B}_A = \mathcal{I}_j(\mathbb{R}^n) \times S^{j-1}$ and hence $\operatorname{Int}\Sigma_A \approx C_{\Gamma/\Gamma_A} \times S^{j-1}$ (see Proposition 5.3). A neighborhood of $\operatorname{Int}\Sigma_A$ in C_Γ is identified with $[0,1) \times C_{\Gamma/\Gamma_A} \times S^{j-1}$ by the homeomorphism onto the image

$$\begin{aligned}&(\varepsilon; f; x_1, \dots, x_{q-1}, x_{q+1}, \dots, x_s; y_{s+1}, \dots, y_{s+t}; v)\\ &\qquad\qquad \longmapsto (f; x_1, \dots, x_{q-1}, x_p + \varepsilon v, x_{q+1}, \dots; y_{s+1}, \dots)\end{aligned} \tag{5.4}$$

where $f \in \mathcal{K}_{n,j}$ and $v \in S^{j-1}$. It is not hard to see that the local coordinate of C_Γ given by (5.4) has the orientation sign $(-1)^{j(q-1)+(j-1)nt}$. Putting $\varepsilon = 0$ in the local coordinate (5.4), we obtain the natural orientation of $C_{\Gamma/\Gamma_A} \times S^{j-1}$. Thus the induced orientation of $C_{\Gamma/\Gamma_A} \times S^{j-1}$ as a boundary face of C_Γ has the sign $(-1)^{j(q-1)+(j-1)nt+1}$.

Case 2: $a = b = 1$ or $a = 0$, $b = 2$.

Let $A = \{x_p, y_q\}$ $(p \leq s < q$ or $s < p < q)$. In this case $\hat{B}_A = \mathcal{I}_j(\mathbb{R}^n) \times S^{n-1}$ and hence $\operatorname{Int}\Sigma_A \approx C_{\Gamma/\Gamma_A} \times S^{n-1}$ (see Proposition 5.3). A neighborhood of $\operatorname{Int}\Sigma_A$ in C_Γ is identified with $[0,1) \times C_{\Gamma/\Gamma_A} \times S^{n-1}$ by the homeomorphism onto the image

$$\begin{aligned}&(\varepsilon; f; x_1, \dots, x_s; y_{s+1}, \dots y_{q-1}, y_{q+1}, \dots, y_{s+t}; w)\\ &\qquad\qquad \longmapsto (f; x_1 \dots; y_{s+1}, \dots, y_{q-1}, z_p + \varepsilon w, y_{q+1}, \dots)\end{aligned} \tag{5.5}$$

where $w \in S^{n-1}$ and $z_p = f(x_p)$ or y_p according to whether p is internal or not. The local coordinate of C_Γ given by (5.5) has the orientation sign $(-1)^{n(s+q-1)+js}$. Putting $\varepsilon = 0$ in the local coordinate (5.5), we obtain the natural orientation of $C_{\Gamma/\Gamma_A} \times S^{n-1}$. Thus the induced orientation of $C_{\Gamma/\Gamma_A} \times S^{n-1}$ as a boundary face of C_Γ has the sign $(-1)^{n(s+q-1)+js+1}$.

5.6. **Principal faces.** In this subsection we compute the fiber integration along the principal faces.

Theorem 5.9 ([5, 14, 20]). *The integration of ω_Γ along the principal face Σ_A of non-infinity type (thus $\sharp A = 2$) vanishes unless the two vertices are joined by an edge in Γ.*

Proof. Let a and b be the numbers of i- and e-vertices in A respectively ($a + b = 2$ since A is principal). If two vertices in A are not connected by an edge, then we have $\hat{\omega}_{\Gamma_A} = 1 \in \Omega^0_{DR}(\hat{B}_A)$. Thus, if A has no i-vertex ($a = 0$, $b = 2$), then the first equality of Lemma 5.4 does not hold;

$$\deg \hat{\omega}_{\Gamma_A} = 0 \neq 0 \cdot j + 2n - (n+1) = n - 1.$$

If A has an i-vertex ($(a,b) = (1,1)$ or $(2,0)$), then the second inequality of Lemma 5.4 does not hold since

$$\deg \hat{\omega}_{\Gamma_A} - (ja + nb - (j+1)) = \begin{cases} -(j-1) & \text{if } (a,b) = (2,0) \\ -(n-1) & \text{if } (a,b) = (1,1). \end{cases}$$

□

Theorem 5.10 ([5, 14, 20]). *Let Γ be an admissible graph. Then the integration of ω_Γ along the principal face Σ^∞_A (thus $\sharp A = 1$) always vanishes.*

Proof. Let p be the only vertex in A. The graph $\Gamma^\infty_A = \Gamma/\Gamma^c_A$ has two vertices; one is p and the other is v^∞. $\sharp E(\Gamma^\infty_A)$ is equal to the valency of p in Γ.

By Lemma 5.7, $(\pi_\Gamma^{\partial_A^\infty})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A^\infty}$ vanishes unless

$$\deg \omega_{\Gamma_A^\infty} = \begin{cases} j-1 & \text{if } p \text{ is internal,} \\ n-1 & \text{if } p \text{ is external,} \end{cases}$$

hence $(\pi_\Gamma^{\partial_A^\infty})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A^\infty}$ does not vanish only if, in Γ,

- p is a uni-valent i-vertex with exactly one adjacent η-edge, or
- p is a uni-valent e-vertex with exactly one adjacent θ-edge.

But neither case occurs since Γ is admissible (see Definition 2.1). □

Theorem 5.11 ([5, 14, 20])**.** *The sum of integrations of ω_Γ along all the principal faces Σ_A of non-infinity type is equal to $I(\delta\Gamma)$.*

Proof. By the above Theorems 5.9 and 5.10, we only need to consider the principal faces Σ_A such that the two vertices of Γ_A are joined by an edge e. We will show

$$(\pi_\Gamma^{\partial_A})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = (-1)^{\tau(e)}(\pi_{\Gamma/e})_*\omega_{\Gamma/e}$$

for any principal strata Σ_A, where $e \in E(\Gamma)$ is the only edge of the subgraph Γ_A and $\tau(e)$ is the sign given in Proposition 2.9. Then we will obtain

$$\sum_{\substack{A\subset V(\Gamma)\\ \sharp A=2}} (\pi_\Gamma^{\partial_A})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = \sum_{e\in E(\Gamma)\setminus\{\text{loops}\}} (-1)^{\tau(e)}(\pi_{\Gamma/e})_*\omega_{\Gamma/e} = I(\delta\Gamma).$$

First we consider the case when n and j are odd. We divide the proof into four cases.

Case (a). $A=\{p,q\}$ consists of two e-vertices (so we can assume $s<p<q$ and the edge $e=\overrightarrow{pq}$ is a θ-edge).

In this case $\hat{B}_A = S^{n-1}$ and $B_A=\{*\}$, so $\mathrm{Int}\,\Sigma_A \approx C^o_{\Gamma/e}\times S^{n-1}$. The induced orientation of $C^o_{\Gamma/e}\times S^{n-1}$ from C_Γ is $(-1)^{n(s+q-1)+js+1}$ by Case 2 in §5.5, and is equal to $(-1)^q$ since n and j are odd. This sign is $(-1)^{\tau(e)}$ (see Proposition 2.9).

Under the identification $\mathrm{Int}\,\Sigma_A \approx C^o_{\Gamma/e}\times S^{n-1}$, the map $\varphi_e : C^o_\Gamma \to S^{n-1}$ restricts to the projection $pr_2 : C^o_{\Gamma/e}\times S^{n-1}\to S^{n-1}$. Hence via the diffeomorphism $\mathrm{Int}\,\Sigma_A \approx S^{n-1}\times C^o_{\Gamma/e}$, the form $\omega_{\Gamma_A}|_{\mathrm{Int}\,\Sigma_A}$ corresponds to

$$(-1)^{\tau(e)}p_A^*\omega_{\Gamma/e}\wedge pr_2^* vol_{S^{n-1}} \in \Omega^*_{DR}(S^{n-1}\times C^o_{\Gamma/e})$$

and hence we have

$$\begin{aligned}(\pi_\Gamma^{\partial_A})_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} &= (-1)^{\tau(e)}(\pi_{\Gamma/e})_*\circ(p_A)_*(p_A^*\omega_{\Gamma/e}\wedge pr_2^* vol_{S^{n-1}}) \\ &= (-1)^{\tau(e)}(\pi_{\Gamma/e})_*\left(\int_{S^{n-1}} vol_{S^{n-1}}\right)\omega_{\Gamma/e} \\ &= (-1)^{\tau(e)}(\pi_{\Gamma/e})_*\omega_{\Gamma/e}.\end{aligned}$$

Case (b). $A=\{p,q\}$ contains both an e- and an i-vertex (thus we can assume $p\le s<q$, and $e=\overrightarrow{pq}$ is a θ-edge).

In this case $\mathrm{Int}\,\Sigma_A \approx C^o_{\Gamma/e}\times S^{n-1}$. Similarly as in Case (a) above, the induced orientation of $C^o_{\Gamma/e}\times S^{n-1}$ from C_Γ is $(-1)^q = (-1)^{\tau(e)}$ and $\varphi_e : C^o_\Gamma \to S^{n-1}$ restricts to the projection $pr_2 : C^o_{\Gamma/e}\times S^{n-1}\to S^{n-1}$. Thus, as in the Case (a),

$$(\pi_\Gamma)_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = (-1)^{\tau(e)}(\pi_{\Gamma/e})_*\omega_{\Gamma/e}.$$

Case (c). Both two points p, q of A are internal and $e = \overrightarrow{pq}$ is an η-edge.

In this case $\mathrm{Int}\,\Sigma_A \approx C^o_{\Gamma/e} \times S^{j-1}$. Proof is the same as the above cases, since by Case 1 in §5.5, the induced orientation of $C^o_{\Gamma/e} \times S^{j-1}$ from C_Γ is $(-1)^q = (-1)^{\tau(e)}$.

Case (d). $A = \{p, q\}$ consists of two i-vertices (thus we can assume $p < q \le s$) and the edge $e = \overrightarrow{pq}$ is a θ-edge.

In this case $\mathrm{Int}\,\Sigma_A \approx C^o_{\tilde{\Gamma}} \times S^{j-1}$, where $\tilde{\Gamma}$ is Γ/e with its small loop e removed. The right hand side is nothing but the space $C^o_{\Gamma/e}$ (Γ/e is a graph with small loop; see §3.3), up to the orientation. Again by Case 1 in §5.5, the induced orientation of $\mathrm{Int}\,\Sigma_A \subset C_\Gamma$ has the sign $(-1)^{\tau(e)}$. Hence

$$(\pi_\Gamma)_* \omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = (-1)^{\tau(e)} (\pi_{\Gamma/e})_* \omega_{\Gamma/e}$$

as desired.

The proof of the case when n and j are even is similar. In Case (a), the induced orientation of $\Sigma_A \subset C_\Gamma$ has the sign -1. To integrate the form θ_e first, we must put θ_e at the top of ω_Γ. Such a re-ordering yields the sign $(-1)^{i-1}$ if e is the i-th edge. Hence

$$(\pi^{\partial_A}_\Gamma)_* \omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = (-1)^i (\pi_{\Gamma/e})_* \omega_{\Gamma/e} = (-1)^{\tau(e)} (\pi_{\Gamma/e})_* \omega_{\Gamma/e}.$$

The remaining three cases are proved in similar ways. In Case (d), we have to put the S^{j-1}-factor at the end of the $(S^{j-1})^u$-part, so the sign $(-1)^u$ appears.

When n is even, we have to consider one more case;

Case (e). $A = \{p, q\}$ consists of two internal vertices (thus we can assume $p < q \le s$) which are joined by an η-edge $\overrightarrow{pq}_\eta$ and a θ-edge $\overrightarrow{pq}_\theta$.

In this case, the induced orientation of $\Sigma_A \subset C_\Gamma$ has the sign -1 as in Case (a). But we need to put S^{j-1} at the end of S^{j-1}-factor, and not to move the forms θ and η. Hence

$$(\pi^{\partial_A}_\Gamma)_* \omega_\Gamma|_{\mathrm{Int}\,\Sigma_A} = (-1)^{u+1} (\pi_{\Gamma/e})_* \omega_{\Gamma/e} = (-1)^{\tau(e)} (\pi_{\Gamma/e})_* \omega_{\Gamma/e}.$$

□

By the above Theorem 5.11, the proof of Theorem 3.4 is reduced to showing that hidden faces do not contribute to the fiber integration. The following, whose proof will be given in §5.7 and §5.8, will complete the proof of Theorem 3.4.

Theorem 5.12. *Let Γ be an admissible graph. Then all the integrations of ω_Γ along hidden boundary faces of the fiber of $\pi_\Gamma : C_\Gamma \to \mathcal{K}_{n,j}$ vanish if (1) $n - j \ge 2$ is even and Γ is a tree, or (2) both $n > j \ge 3$ are odd and Γ has at most one loop component.*

5.7. **Hidden faces; the non-infinity type.**

Here we show that all the hidden faces Σ_A of non-infinity type, $\sharp A \ge 3$, do not contribute to the fiber integration.

Lemma 5.13 ([14, 20]). *Suppose $A \subset V(\Gamma)$ is such that the subgraph Γ_A is not connected. Then $(\pi^{\partial_A}_\Gamma)_* \omega_\Gamma|_{\mathrm{Int}\,\Sigma_A}$ vanishes.*

Proof. If Σ_A is principal (so $\sharp A = 2$), then the claim of this Lemma is exactly that of Theorem 5.9. So we assume $\sharp A \ge 3$.

Suppose $\Gamma_A = \Gamma_{A_1} \sqcup \Gamma_{A_2}$ for non-empty subsets $A_1, A_2 \subset A$ (thus we can assume $\sharp A_1 \ge 2$). Let a_i and b_i be the numbers of i- and e-vertices of A_i, $i = 1, 2$.

We define the space $\tilde{B}_A$, which contains $\hat{B}_A$ as an open subset, by

$$\tilde{B}_A := (C^0_{a_1,b_1} \times C^o_{a_2,b_2}) / \sim$$

here $\sim$ is defined by using the translation and the scaling. In other words, 'a point in A_1 may collide with a point in A_2.' Since there are no edges joining a point in A_1 to that of A_2, the direction maps φ corresponding to the edges of Γ_A are well-defined on $\tilde{B}_A$, and so is the associated differential form (we denote it by $\tilde{\omega}_{\Gamma_A}$). The restriction of $\tilde{\omega}_{\Gamma_A}$ onto $\hat{B}_A$ is $\hat{\omega}_{\Gamma_A}$.

Consider a free action of $\mathbb{R}^N$ on $\tilde{B}_A$ defined by the translations of points in A_1 (points in A_2 are fixed). Here $N = j$ or n according to whether A_1 contains an i-vertex or not. Let $p : \tilde{B}_A \to \tilde{B}_A/\mathbb{R}^N$ be the quotient map. Since the direction maps $\tilde{\varphi} : \tilde{B}_A \to S^{j-1}$ or S^{n-1} factor through p, there exists a form $\omega'_{\Gamma_A} \in \Omega^*_{DR}(\tilde{B}_A/\mathbb{R}^N)$ such that $p^*\omega'_{\Gamma_A} = \tilde{\omega}_{\Gamma_A}$. We have a map of fibrations

$$\begin{tikzcd}
\hat{B}_A \arrow[r, hook] \arrow[dr, "\rho_A"'] & \tilde{B}_A \arrow[r, two heads, "p"] \arrow[d, "\tilde{\rho}_A"] & \tilde{B}_A/\mathbb{R}^N \arrow[dl, "\rho'_A"] \\
 & B_A &
\end{tikzcd}$$

(for definition of ρ_A see Proposition 5.3) and it holds that

$$(\rho_A)_*\omega_{\Gamma_A} = (\tilde{\rho}_A)_*\tilde{\omega}_{\Gamma_A} = (\rho'_A)_*\omega'_{\Gamma_A}.$$

This implies $(\rho_A)_*\omega_{\Gamma_A} = 0$, since the fiber of ρ'_A is of strictly less dimension than those of ρ_A and $\tilde{\rho}_A$. This together with the formula (5.3) completes the proof. $\square$

Thanks to Lemma 5.13, below we can assume that Γ_A is connected.

In Theorem 3.4 we assumed $g \leq 1$, that is, our graph has at most one loop component (other than small loops), and so does its connected subgraph Γ_A.

Proposition 5.14. *Suppose $n - j$ is even. If $\sharp A \geq 3$ and Γ_A is a tree, then $(\pi^{\partial_A}_{\Gamma})_*\omega_{\Gamma}|_{\mathrm{Int}\,\Sigma_A}$ vanishes.*

Proof. Since Γ_A is a tree, there are at least two uni-valent vertices in Γ_A. All the possibilities of uni-valent vertices are listed in Figure 5.3. We will prove the vanishing of integration along Σ_A in the successive Lemmas; types (a) and (c) in Lemma 5.15 and type (b) in Lemma 5.16. The assumption that $n - j$ is even will be used in the proof of Lemma 5.16. $\square$



FIGURE 5.3. Uni-valent vertices

Lemma 5.15 ([5])**.** *If $\sharp A \geq 3$ and A has a uni-valent vertex p of types (a) or (c) in Figure 5.3, then the integration $(\pi^{\partial_A}_{\Gamma})_*\omega_{\Gamma}|_{\mathrm{Int}\,\Sigma_A}$ vanishes.*

Proof. Let $q \in A$ be the vertex joined to p in Γ_A, which must be internal in the case (a), while in the case (c) it may be both internal or external. There is a fiberwise free action $\mathbb{R}_{>0}$ on $\hat{B}_A$ defined on each fiber by

$$(\dots, z_p, \dots) \longmapsto (\dots, az_p + (1-a)z_q, \dots), \quad a \in \mathbb{R}_{>0}.$$

Then $\hat{\omega}_{\Gamma_A} \in \Omega^*_{DR}(\hat{B}_A)$ is basic with respect to the quotient $\hat{B}_A \to \hat{B}_A/\mathbb{R}_{>0}$. Since $\sharp A \geq 3$, the fiber of $\hat{B}_A/\mathbb{R}_{>0} \to B_A$ is of strictly less dimension than that of ρ_A. Hence the similar argument as in Lemma 5.13 completes the proof. $\square$

Lemma 5.16. *Suppose $n-j$ is even. If $\sharp A \geq 3$ and Γ_A is a tree all of whose uni-valent vertices are of type (b) in Figure 5.3, then $(\pi_\Gamma)_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A}$ vanishes.*

Proof. The vertex q of A which is joined to p may be both internal or external. Since valency of q is greater than one, there exist vertices $r_1, \dots, r_a$ ($r_i \neq p$, $a \geq 1$) which are joined to q.

Suppose one of them, say r_1, is uni-valent. By our assumption r_1 is also of type (b). Consider a fiberwise involution $\chi_1 : \hat{B}_A \to \hat{B}_A$ defined by

$$\chi_1(\iota; \dots, x_p, \dots, x_{r_1}, \dots) := (\iota; \dots, x_{r_1}, \dots, x_p, \dots)$$

(other coordinates are not changed). This involution changes the orientation of the fiber by $(-1)^j$, while $\chi_1^*\hat{\omega}_{\Gamma_A} = (-1)^{n-1}\hat{\omega}_{\Gamma_A}$ since $\chi_1^*\theta_{pq} = \theta_{r_1 q}$, $\chi_1^*\theta_{r_1 q} = \theta_{pq}$ and θ's are of degree $n-1$. Thus

$$(\rho_A)_*\hat{\omega}_{\Gamma_A} = (-1)^{j+n-1}(\rho_A)_*\hat{\omega}_{\Gamma_A} = -(\rho_A)_*\hat{\omega}_{\Gamma_A}$$

since $n-j$ is even, and it must vanish.

Thus we may assume all the r_i's are at least bi-valent. That is, we can assume that all the uni-valent vertices p (of type (b)) and adjacent q are such that no other uni-valent vertex is joined to q. Then we can find at least two pairs (p, q) of vertices such that p is uni-valent vertex joined to exactly one bi-valent vertex q; since otherwise Γ_A cannot be a tree. Such pairs, say (p, q) and (p', q'), are of types (b-1) or (b-2) or (b-3) in Figure 5.4, where an asterisk can be both internal or external.



FIGURE 5.4. Types (b-1), (b-2), (b-3)

If (p, q) is of type (b-1), define a fiberwise involution $\chi_2 : \hat{B}_A \to \hat{B}_A$ by

$$\chi_2(\iota; \dots, y_q, \dots) := (\iota; \dots, \iota(x_p) + z_r - y_q, \dots),$$

where z_r is $\iota(x_r)$ or y_r according to whether r is internal or not. This involution has orientation sign $(-1)^n$, while $\chi_2^*\hat{\omega}_{\Gamma_A} = (-1)^{n-1}\hat{\omega}_{\Gamma_A}$ similarly to χ_1. Thus

$$(\rho_A)_*\hat{\omega}_{\Gamma_A} = (-1)^{n+n-1}(\rho_A)_*\hat{\omega}_{\Gamma_A} = -(\rho_A)_*\hat{\omega}_{\Gamma_A}$$

and it must vanish.

So finally we can assume that both pairs (p, q) and (p', q') are of types (b-2) or (b-3). The proof of this case appeared in [14, 20]; consider a fiberwise involution $\chi_3 : \hat{B}_A \to \hat{B}_A$ defined by

$$\chi_3(\iota; \dots, x_p, \dots, x_{p'}, \dots) := (\iota; \dots, x_q - x_{q'} + x_{p'}, \dots, x_q - x_{q'} + x_p, \dots).$$

The orientation sign of χ_3 is $(-1)^j$, and it satisfies $\chi_3^*\hat{\omega}_{\Gamma_A} = (-1)^{n-1}\hat{\omega}_{\Gamma_A}$ since $\chi_3^*\theta_{pq} = \theta_{p'q'}$, $\chi_3^*\theta_{p'q'} = \theta_{pq}$. Since $n-j$ is even,

$$(\rho_A)_*\hat{\omega}_{\Gamma_A} = (-1)^{n+j-1}(\rho_A)_*\hat{\omega}_{\Gamma_A} = -(\rho_A)_*\hat{\omega}_{\Gamma_A}$$

as in the case of χ_1, and it must vanish. □

When both $n > j \geq 3$ are odd, we have to consider the case that Γ_A has one loop component to prove Theorem 3.4.

Proposition 5.17 ([14, 20])**.** *Suppose $n > j \geq 3$ are odd. If $\sharp A \geq 3$ and Γ_A is connected with one loop component, then $(\pi_\Gamma)_*\omega_\Gamma|_{\mathrm{Int}\,\Sigma_A}$ vanishes.*

Proof. Let s and t be the numbers of i- and e-vertices in A respectively. Define an involution F of $\hat{B}_A$ by

$$F(\iota; x_1, \dots, x_s; y_{s+1}, \dots, y_{s+t}) := \\ (\iota; x_1, 2x_1 - x_2, \dots, 2x_1 - x_s; 2\iota(x_1) - y_{s+1}, \dots, 2\iota(x_1) - y_{s+t}),$$

whose orientation sign is $(-1)^{j(s-1)+nt} = (-1)^{s+t-1}$.

Let α and β be the numbers of η- and θ-edges of Γ_A respectively. Since $\hat{\varphi}_e \circ F = \iota_{S^{N-1}} \circ \hat{\varphi}_e$ for any edge e ($N = j$ or n according to whether e is an η-edge or a θ-edge), we have

$$F^*\omega_{\Gamma_A} = (-1)^{j\alpha + n\beta}\omega_{\Gamma_A} = (-1)^{\alpha+\beta}\omega_{\Gamma_A}.$$

But by our assumption, Γ_A has exactly one loop component, so $\alpha + \beta$ is equal to the number of vertices of Γ_A, that is, $s + t$. Thus

$$(\rho_A)_*\omega_{\Gamma_A} = (-1)^{s+t-1}(-1)^{s+t}(\rho_A)_*\omega_{\Gamma_A} = -(\rho_A)_*\omega_{\Gamma_A}$$

and hence $(\rho_A)_*\omega_{\Gamma_A} = 0$. The formula (5.3) in §5.3 completes the proof. □

5.8. **Hidden faces; strata at infinity.**

In this subsection we will prove that the hidden strata Σ_A^∞ at infinity (thus $\sharp A \geq 2$) do not contribute to integrals.

Lemma 5.18 ([5, 14, 20])**.** *If $\sharp A \geq 2$, then the integration along Σ_A^∞ vanishes.*

Proof. First we will show that $\omega_{\Gamma_A^\infty}$ cannot satisfy the equation in Lemma 5.7 when $A \subset V(\Gamma)$ is a proper subset.

If Γ is an admissible graph, then each e-vertex of Γ_A^∞ ($= \Gamma/\Gamma_A^c$) is of at least tri-valent, and each i-vertex is an endpoint of some θ-edge. This implies

$$2\sharp\{\theta\text{-edges of } \Gamma_A^\infty\} \geq a + 3b, \tag{5.6}$$

where a and b are numbers of i- and e-vertices of Γ_A.

Since $\Gamma_A^\infty \setminus \{v^\infty\}$ is a one-dimensional open object with at most one loop component and with at least one open edge,

$$\sharp\{\eta\text{-edges of } \Gamma_A^\infty\} + \sharp\{\theta\text{-edges of } \Gamma_A^\infty\} \geq a + b. \tag{5.7}$$

By using estimations (5.6), (5.7) and $n - j - 2 \geq 0$, we see that $\omega_{\Gamma_A^\infty}$ does not satisfy the criterion of Lemma 5.7;

$$\begin{aligned}
\deg \omega_{\Gamma_A^\infty} &\geq (n-1)\sharp\{\theta\text{-edges of } \Gamma_A^\infty\} + (j-1)(a + b - \sharp\{\theta\text{-edges of } \Gamma_A^\infty\}) \\
&= (n-j)\sharp\{\theta\text{-edges of } \Gamma_A^\infty\} + (j-1)(a+b) \\
&\geq \frac{1}{2}(n-j)(a+3b) + (j-1)(a+b) \\
&= (ja + nb - 1) + \frac{1}{2}(n-j-2)(a+b) + 1 \\
&> ja + nb - 1 \\
&= \dim \hat{B}_A^\infty.
\end{aligned}$$

Next consider the case $A = V(\Gamma)$ (thus $\Gamma_A^\infty = \Gamma$). Define a fiberwise free action of $\mathbb{R}_{>0}$ on $\hat{B}_A^\infty$ by

$$(x_1, \dots, y_{a+1}, \dots) \longmapsto (x_1, \alpha x_2 + (1-\alpha)x_1, \dots, \alpha y_{a+1} + (1-\alpha)f_0(x_1), \dots),$$

where $f_0 : \mathbb{R}^j \hookrightarrow \mathbb{R}^n$ is the standard inclusion given by $x \mapsto (x, 0, \dots, 0)$. This action is non-trivial since $\sharp A \geq 2$. The differential form $\omega_{\Gamma^\infty_{V(\Gamma)}}$ is basic with respect to this action, hence similar argument to Lemma 5.13 completes the proof. □

Remark 5.19. In the end of the proof we used that the long knots are standard near infinity, so $\omega_{\Gamma^\infty_{V(\Gamma)}}$ contains no information about the base space $\mathcal{K}_{n,j}$.

Thus we have shown that all of hidden and infinity contributions vanish, and completed the proof of Theorem 5.12.

5.9. **Proof of Theorem 4.1.** In the above proofs, we have used

- symmetry of the fiber, and
- dimension counting.

Below we check that some of the arguments are valid even if $n - j \geq 3$ is odd (in particular $n = 6k$, $j = 4k - 1$), and can be used to prove Theorem 4.1.

Lemma 5.20. *$d\mathcal{H} = I(\delta H)$ modulo the contributions of hidden faces.*

Proof. Exactly similar to Theorem 5.11. We used Theorems 5.9 and 5.10 to prove Theorem 5.11, which are proved by only dimension counting. □

Lemma 5.21. *The hidden faces except for the face Σ_A corresponding to $A = V(H_2)$ do not contribute to the integral.*

Proof. The strata at infinity do not contribute; the proof of Lemma 5.18 does not use the parities of n, j. By Lemma 5.13, which uses only dimension counting, we have only to consider the faces Σ_A for A such that the subgraph Γ_A is connected.

For $A \subset V(H_1)$ with $\sharp A = 3$, use Lemma 5.15.

For $A = V(H_1)$, consider a fiberwise involution $F : \hat{B}_A \to \hat{B}_A$ defined by

$$F(\iota; (x_1, \dots, x_4)) := (\iota; (2x_2 - x_1, x_2, x_3, x_4)).$$

The orientation sign of F is $(-1)^j$, while $F^*\hat{\omega}_{H_1} = (-1)^n\hat{\omega}_{H_1}$ because $F^*\theta_{12} = (-1)^n\theta_{12}$ and F^* preserves η_{23}, θ_{34}. Since $n - j$ is odd,

$$(\rho_A)_*\hat{\omega}_{H_1} = (-1)^{n+j}(\rho_A)_*\hat{\omega}_{H_1} = -(\rho_A)_*\hat{\omega}_{H_1}$$

and hence the integration along $\Sigma_{V(H_1)}$ must vanish.

For $A \subset V(H_2)$ with $\sharp A = 3$, then A contains two i-vertices joined to the e-vertex labeled by 4. Then using the involution χ_2 appeared in the proof of Lemma 5.16, vanishing for type (b-1), we can complete the proof. □

Next consider the contribution of 'anomalous face' Σ_A, $A = V(H_2)$ (hence $\Gamma_A = H_2$). Recall from §5.3 that the face Σ_A is described by the pullback square in Proposition 5.3 with $B_A = \mathcal{I}_j(\mathbb{R}^n)$ and $C_{\Gamma/\Gamma_A} = \mathcal{K}_{n,j} \times \mathbb{R}^j$. The contribution of Σ_A is given by $p_*D_A^*(\rho_A)_*\hat{\omega}_{H_2}$, where $p : \mathcal{K}_{n,j} \times \mathbb{R}^j \to \mathcal{K}_{n,j}$ is the first projection.

At present we cannot determine whether this contribution vanishes or not, so

$$dI(H) = \frac{1}{6}p_*D_A^*(\rho_A)_*\hat{\omega}_{\Gamma_A}.$$

But we can define a correction term c which kills this contribution as follows.

Lemma 5.22. *The form $(\rho_A)_*\hat{\omega}_{H_2} \in \Omega^{2n-2j-2}_{DR}(\mathcal{I}_j(\mathbb{R}^n))$ is closed.*

Proof. By the generalized Stokes theorem,

$$d(\rho_A)_*\hat{\omega}_{H_2} = \pm(\rho_A^\partial)_*\hat{\omega}_{H_2},$$

where ρ_A^∂ is the restriction of ρ_A onto the boundary of the fiber. The principal faces correspond to the graph (up to labeling)



obtained by contracting the edge $i4$ ($i = 1, 2, 3$) of H_2. Consider the involution

$$F : (x_1, x_2, x_3) \longmapsto (x_1, x_2, 2x_2 - x_3)$$

of the principal face. The orientation sign of F is $(-1)^j$, while $F^*\hat{\omega}_{H_2} = (-1)^n\hat{\omega}_{H_2}$ since $F^*\theta_{12} = \theta_{12}$ and $F_1^*\theta_{23} = \theta_{32} = (-1)^n\theta_{23}$. Since $j + n$ is odd, the integration along the principal faces vanish.

The hidden (but non-anomalous) contributions are proved to vanish in completely similar way as in Lemma 5.21. Since there is no anomalous face of the fiber of $\hat{B}_A \to B_A$, we have proved that $d(\rho_A)_*\hat{\omega}_{H_2} = \pm(\rho_A^\partial)_*\hat{\omega}_{H_2} = 0$. □

Thus we have a cohomology class $[(\rho_A)_*\hat{\omega}_{H_2}] \in H_{DR}^{2n-2j-2}(\mathcal{I}_j(\mathbb{R}^n))$. But in fact $\mathcal{I}_j(\mathbb{R}^n)$ is homotopy equivalent to Stiefel manifold of j-frames in $\mathbb{R}^n$ [14], and it is known that, when $n - j$ is odd, its cohomology ring with coefficients in $\mathbb{R}$ is given by

$$H^*(\mathcal{I}_j(\mathbb{R}^n); \mathbb{R}) \cong \begin{cases} H^*(S^{2n-5} \times S^{2n-9} \times \cdots \times S^{2(n-j)+1} \times S^{n-1}; \mathbb{R}) & n \text{ is even,} \\ H^*(S^{2n-7} \times S^{2n-11} \times \cdots \times S^{2(n-j)+1}; \mathbb{R}) & n \text{ is odd.} \end{cases}$$

Hence we can find a form $\mu \in \Omega_{DR}^{2n-2j-3}(\mathcal{I}_j(\mathbb{R}^n))$ such that $d\mu = (\rho_A)_*\hat{\omega}_{H_2}/6$ (the factor S^{n-1} in the right hand side does not cause any trouble; if $2n-2j-2 = n-1$ then $n = 2j+1$ and n becomes odd).

Definition 5.23. Define $c := -p_*D_A^*\mu \in \Omega_{DR}^{2n-3j-3}(\mathcal{K}_{n,j})$ ($p : \mathcal{K}_{n,j} \times \mathbb{R}^j \to \mathcal{K}_{n,j}$ is the first projection).

Then Theorem 4.1 is easily proved;

$$\begin{aligned} d(I(H) + c) &= p_*D_A^*(\rho_A)_*\hat{\omega}_{H_2}/6 - dp_*D_A^*\mu \\ &= p_*D_A^*(\rho_A)_*\hat{\omega}_{H_2}/6 - p_*dD_A^*\mu \pm p_*^\partial D_A^*\mu \qquad &&\text{(Stokes theorem)} \\ &= p_*D_A^*(\rho_A)_*\hat{\omega}_{H_2}/6 - p_*D_A^*d\mu &&(D_A \text{ is constant near } \partial C_1(\mathbb{R}^j)) \\ &= 0 &&\text{(by definition of } \mu). \end{aligned}$$

Remark 5.24. We considered involutions $i_{p,q} : \mathbb{R}^n \to \mathbb{R}^n$ in the proof of Theorem 4.19. They induce involutions $i_{p,q}$ on $\mathcal{I}_j(\mathbb{R}^n)$ given by $\iota \mapsto i_{p,q} \circ \iota$. This lifts to $\hat{i}_{p,q} : \hat{B}_A \to \hat{B}_A$ ($A = V(H_2)$) defined by $\hat{i}_1(\iota; (x_1, x_2, x_3); y) := (i_{p,q} \circ \iota; (x_1, x_2, x_3); i_{p,q}(y))$. This has the orientation sign $(-1)^{p-q+1}$ on the fiber, and $\hat{i}_1^*\hat{\omega}_A = (-1)^{p-q+1}\hat{\omega}_A$ since $i_{p,q}^*\theta_{*4} = (-1)^{p-q+1}\theta_{*4}$ (see the remark after Proposition 4.2). Hence $i_{p,q}^*(\rho_A)_*\hat{\omega}_A = (\rho_A)_*\hat{\omega}_A$. This implies that, replacing μ with $(\mu + i_{p,q}^*\mu)/2$, we may assume that $i_{p,q}^*\mu = \mu$. Since $i_{p,q}$'s for different p, q commute with each other, we can arrange μ so that it is preserved by all of these involutions by repeating the same procedure as above.

5.10. **Independency on volume forms.** There is another vanishing result, which is needed in the proof of Propositions 3.5, 3.6 (independency on the choices of volume forms of the map I on cohomology). Recall that we assume that g, n and j are such that the integration map I is a cochain map and that $n-j>2$.

Lemma 5.25. *The differential form $\tilde{I}(\Gamma)\in\Omega^*_{DR}(\mathcal{K}_{n,j}\times[0,1])$ from the proof of Proposition 3.5 is closed if $n-j>2$.*

Proof. As already done for $I(\Gamma)$ in §5.6, §5.7 and §5.8, we must show

- $d_{\mathcal{K}_{n,j}\times[0,1]}\tilde{I}(\Gamma)=\tilde{I}(\delta\Gamma)$ modulo the contributions of hidden faces of the fibers of π_{Γ_i}, and hence it vanishes since Γ is a cocycle, and
- the contributions of hidden faces also vanish.

In this section we have proved the vanishing results by using symmetry and dimension counting. We have to repeat these proofs for $\tilde{I}$. The symmetry arguments can be applied to the cases here, since the factor $[0,1]$ does not cause any trouble.

We can check that the dimension-counting arguments also work. The key ingredients in the proofs are the equation (5.3) and Lemmas 5.4, 5.7. For $\tilde{I}$, we need to replace the criteria in Lemmas 5.4, 5.7 with

$$\begin{array}{ll}\deg\omega_{\Gamma_A}=nb-n \ \text{ or } \ nb-(n+1) & \text{if } a=0,\\ 0\le\deg\omega_{\Gamma_A}-(ja+nb-(j+1))\le nj+1 & \text{if } a>0.\end{array}$$

and

$$\deg\omega_{\Gamma_A^\infty}=ja+nb \ \text{ or } \ ja+nb-1$$

respectively, since these criteria are consequences of the following pullback square

$$\begin{tikzcd}
\operatorname{Int}\Sigma_A\times[0,1] \arrow[r, "\hat{D}_A\times\mathrm{id}"] \arrow[d, "p_A\times\mathrm{id}"'] & \hat{B}_A\times[0,1] \arrow[d, "\rho_A\times\mathrm{id}"] \\
C^o_{\Gamma/\Gamma_A}\times[0,1] \arrow[r, "D_A\times\mathrm{id}"] & B_A\times[0,1]
\end{tikzcd}$$

and of $\operatorname{Int}\Sigma_A^\infty\times[0,1]\approx C_{\Gamma_A^c}\times\hat{B}_A^\infty\times[0,1]$. These replacements do not affect all the arguments in §5.7 and §5.8, except for Lemma 5.18; a problem may occur when $n-j=2$ and $A=\{p,q\}$ consists of two i-vertices, since in such a case $\deg\omega_{\Gamma_A^\infty}=(j-1)+(n-1)=2j=\dim(\hat{B}_A^\infty\times[0,1])$ can happen. $\square$

Similar arguments show the following.

Lemma 5.26. *Suppose $n-j\ge3$ is odd. Then the cohomology class $\mathcal{H}\in H^{2n-3j-3}_{DR}(\mathcal{K}_{n,j})$ is independent of the choices of (anti-)symmetric volume forms.*

Proof. First let $vol_{S^{j-1}}$ be fixed, and let v_0, v_1 be two symmetric volume forms of S^{n-1} with total integral one. Define the maps I_0, I_1, $\tilde{I}$ and the form $\tilde{v}$ as in the proof of Proposition 3.5. Then we have

- $d_{\mathcal{K}_{n,j}\times[0,1]}\tilde{I}(H)=\tilde{I}(\delta H)$ modulo the contributions of hidden faces of the fibers of π_{H_i}, and hence vanishes since H is a cocycle, and
- the contributions of hidden faces except for $\Sigma_{V(H_2)}$ also vanish.

Let $A:=V(H_2)$. Then $(\rho_A\times\mathrm{id})_*\hat{\omega}_A\in\Omega^{2n-2j-2}_{DR}(\mathcal{I}_j(\mathbb{R}^n)\times[0,1])$ is a closed form; the proof is same as Lemma 5.22. But since $H^{2n-2j-2}_{DR}(\mathcal{I}_j(\mathbb{R}^n)\times[0,1])=0$ when $n-j$ is odd, we have a form $\tilde{\mu}$ which satisfies $(\rho_A\times\mathrm{id})_*\hat{\omega}_A/6=d\tilde{\mu}$. Using

$\tilde{c} := -\tilde{p}_*(D_A \times \mathrm{id})^*\tilde{\mu}$ (where $\tilde{p} : \mathbb{R}^j \times \mathcal{K}_{n,j} \times [0,1] \to \mathcal{K}_{n,j} \times [0,1]$ is the projection), we have a closed form

$$\tilde{\mathcal{H}} := \tilde{I}(H) + \tilde{c} \in \Omega_{DR}^{2n-3j-3}(\mathcal{K}_{n,j} \times [0,1]).$$

By the generalized Stokes theorem,

$$(I_1(H) + \tilde{c}|_{\mathcal{K}_{n,j}\times\{1\}}) - (I_0(H) + \tilde{c}|_{\mathcal{K}_{n,j}\times\{0\}}) = \pm dp_*\tilde{\mathcal{H}}.$$

But $\tilde{c}|_{\mathcal{K}_{n,j}\times\{\varepsilon\}}$ ($\varepsilon = 0,1$) comes from $\tilde{\mu}|_{\mathcal{I}_j(\mathbb{R}^n)\times\{\varepsilon\}}$ which satisfies $d\tilde{\mu}|_{\mathcal{I}_j(\mathbb{R}^n)\times\{\varepsilon\}} = (\rho_A \times \mathrm{id}_{\{\varepsilon\}})_*\hat{\omega}_A/6$. Hence $\tilde{c}|_{\mathcal{K}_{n,j}\times\{\varepsilon\}}$ works as a correction term for $I_\varepsilon(H)$. Thus we see that $\mathcal{H}$ is independent of the choices of symmetric $vol_{S^{n-1}}$.

Similar arguments work when we fix $vol_{S^{n-1}}$ and use two different $vol_{S^{j-1}}$'s (in this case we can choose $\tilde{\mu}$ of the form $q^*\mu$ where $q : \mathcal{I}_j(\mathbb{R}^n) \times [0,1] \to \mathcal{I}_j(\mathbb{R}^n)$ is the projection, because the graph H_2 contains no η-edges and hence we do not use $vol_{S^{j-1}}$ to define the correction term). □

References

[1] D. Altschuler and L. Freidel, *Vassiliev knot invariants and Chern-Simons perturbation theory to all orders*, Comm. Math. Phys. **187** (1997), no. 2, 261–287.

[2] S. Axelrod and I. M. Singer, *Chern-Simons perturbation theory II*, J. Diff. Geom. **39** (1994), 173–213.

[3] R. Bott and C. Taubes, *On the self-linking of knots*, J. Math. Phys. **35** (1994), no. 10, 5247–5287.

[4] R. Budney, *A family of embedding spaces*, Geom. Topol. Monogr. **13** (2008), 41–84.

[5] A. Cattaneo, P. Cotta-Ramusino, and R. Longoni, *Configuration spaces and Vassiliev classes in any dimensions*, Algebr. Geom. Topol. **2** (2002), 949–1000.

[6] A. Cattaneo and C. Rossi, *Wilson surfaces and higher dimensional knot invariants*, Comm. Math. Phys. **256** (2005), no. 3, 513–537.

[7] K. Habiro, T. Kanenobu and A. Shima, *Finite type invariants of ribbon 2-knots*, Contemp. Math. vol. 233, 187–196.

[8] A. Haefliger, *Knotted $(4k-1)$-spheres in $6k$-space*, Ann. of Math. (2) **75** (1962), 452–466.

[9] ———, *Differential embeddings of S^n in S^{n+q} for $q > 2$*, Ann. of Math. (2) **83** (1966), 402–436.

[10] T. Kohno, *Vassiliev invariants and de Rham complex on the space of knots*, Contemp. Math. vol. 179, 123–138.

[11] R. Longoni, *Nontrivial classes in $H^*(Imb\,(S^1,\mathbb{R}^n))$ from nontrivalent graph cocycles*, Int. J. Geom. Methods Mod. Phys. **1** (2004), no. 5, 639–650.

[12] D. Roseman, *Spinning knots about submanifolds; spinning knots about projections of knots*, Topology Appl. **31** (1989), no. 3, 225–241.

[13] D. Roseman and M. Takase, *High-codimensional knots spun about manifolds*, Algebr. Geom. Topol. **7** (2007), 359–377.

[14] C. Rossi, *Invariants of higher-dimensional knots and topological quantum field theories*, Ph.D. thesis, Zurich University, 2002.

[15] K. Sakai, *Nontrivalent graph cocycle and cohomology of the long knot space*, Algebr. Geom. Topol. **8** (2008), 1499–1522.

[16] K. Sakai and T. Watanabe, *1-loop graphs and configuration space integral for embedding spaces*, in preparation, draft available at http://www.ms.u-tokyo.ac.jp/~ksakai/index.html

[17] D. Sinha, *Operads and knot spaces*, J. Amer. Math. Soc., **19**, no. 2 (2006), 461–486.

[18] V. Tourtchine (Turchin), *On the other side of the bialgebra of chord diagrams*, J. Knot Theory Ramifications **16** (2007), no. 5, 575–629.

[19] V. Vassiliev, *Combinatorial formulas for cohomology of knot spaces*, Mosc. Math. J. **1** (2001), no. 1, 91–123.

[20] T. Watanabe, *Configuration space integral for long n-knots and the Alexander polynomial*, Algebr. Geom. Topol. **7** (2007), 47–92.

Graduate School of Mathematical Sciences, University of Tokyo
E-mail address: `ksakai@ms.u-tokyo.ac.jp`
URL: `http://www.ms.u-tokyo.ac.jp/~ksakai/index.html`